


Insuffisance Cardiaque Avancée

Dr Maxime FAURE
Service de Traitement de l'Insuffisance Cardiaque
27/04/2023



Insuffisance cardiaque : définition

3.1 Definition of heart failure

Heart failure is not a single pathological diagnosis, but a **clinical syndrome** consisting of cardinal symptoms (e.g. breathlessness, ankle swelling, and fatigue) that may be accompanied by signs (e.g. elevated jugular venous pressure, pulmonary crackles, and peripheral oedema). It is due to a **structural and/or functional abnormality of the heart** that results in **elevated intracardiac pressures and/or inadequate cardiac output at rest and/or during exercise**.

Table 3 Definition of heart failure with reduced ejection fraction, mildly reduced ejection fraction and preserved ejection fraction

Criteria	HFrEF	HFmrEF	HFpEF
1	Symptoms ± Signs ^a	Symptoms ± Signs ^a	Symptoms ± Signs ^a
2	LVEF ≤40%	LVEF 41–49% ^b	LVEF ≥50%
3	—	—	Objective evidence of cardiac structural and/or functional abnormalities consistent with the presence of LV diastolic dysfunction/raised LV filling pressures, including raised natriuretic peptides ^c


Etiologies

Table 1 Causes of heart failure, common modes of presentation and specific investigations


Causes	Common modes of presentation	Specific investigations
Coronary artery disease	Angina or "silent ischaemia"	ECG, stress echocardiography, CT coronary angiography, invasive coronary angiography
Hypertension	Heart failure with preserved systolic function, left ventricular hypertrophy, left atrial enlargement	24h ambulatory BP, echocardiography, renal artery imaging, renal and abdominal aorta - noninvasive imaging
Valve disease	Primary valve disease (e.g. aortic stenosis), secondary valve disease (e.g. functional mitral regurgitation)	Echocardiography, cardiac MRI, aortic valve area, aortic regurgitation, aortic valve calcification
Arrhythmias	Atrial tachyarrhythmias, ventricular arrhythmias	Ambulatory ECG monitoring, electrophysiology study, if indicated
CHF	Left ventricular failure, right ventricular failure	ECG, echocardiography, chest X-ray, BNP, natriuretic peptide testing
Diabetes	Heart failure with preserved systolic function	Echocardiography, HbA1c, renal function, lipid profile
Alcohol	Heart failure with preserved systolic function	Echocardiography, liver function tests, renal function, lipid profile
Ischaemic	Heart failure with reduced systolic function	Echocardiography, CT coronary angiography, invasive coronary angiography
Drug-induced	Heart failure with preserved systolic function	Echocardiography, renal function, liver function tests, renal function, lipid profile
Infectious	Heart failure with preserved systolic function	Echocardiography, blood cultures, renal function, liver function tests, renal function, lipid profile
Storage disorders	Heart failure with preserved systolic function	Echocardiography, renal function, liver function tests, renal function, lipid profile
Endocrine disorders	Heart failure with preserved systolic function	Echocardiography, renal function, liver function tests, renal function, lipid profile
Autoimmune	Heart failure with preserved systolic function	Echocardiography, renal function, liver function tests, renal function, lipid profile
Neurological	Heart failure with preserved systolic function	Echocardiography, renal function, liver function tests, renal function, lipid profile
Genetic	Heart failure with preserved systolic function	Echocardiography, renal function, liver function tests, renal function, lipid profile

Pronostic de l'IC

Étude écoscaise sur 1,75 millions personnes; 56658 pts étudiés



Evolution naturelle de l'IC



Allen LA et al. Circulation.
2012 Apr 17;126(16):1928-52

IC Avancée : définition

Table 13 Criteria for definition of advanced heart failure

- All the following criteria must be present despite optimal medical treatment:**
- Severe and persistent symptoms of heart failure (NYHA class III (advanced) or IV).
 - Severe cardiac dysfunction defined by at least one of the following:
 - LVEF ≤30%
 - Isolated RV failure (e.g., ARVC)
 - Non-operable severe valve abnormalities
 - Non-operable severe congenital abnormalities
 - Persistently high (or increasing) BNP or NT-proBNP values and severe LV diastolic dysfunction or structural abnormalities (according to the definitions of HFpEF).
 - Episodes of pulmonary or systemic congestion requiring high-dose i.v. diuretics (or diuretic combinations) or episodes of low output requiring inotropes or vasoactive drugs or malignant arrhythmias causing >1 unplanned visit or hospitalization in the last 12 months.
 - Severe impairment of exercise capacity with inability to exercise or low 6MWT distance (<300 m) or pVO₂ <12 mL/kg/min or <50% predicted value, estimated to be of cardiac origin.

The updated HFA-ESC 2018 criteria for the definition of advanced HF are reported in Table 13.^{37b} A severely reduced LVEF is common but not required for a diagnosis of advanced HF as it may develop in patients with HFpEF as well. In addition to the reported criteria, extra-cardiac organ dysfunction due to HF (e.g. cardiac cachexia, liver or kidney dysfunction) or type II pulmonary hypertension may be present, but are not required for the definition of advanced HF.^{37b}

IC avancée: marqueurs biologiques


10. Advanced heart failure

Supplementary Table 13 Suggested clinical, laboratory, and echocardiographic criteria to trigger referral to a specialized heart failure or advanced heart failure unit

Clinical	Laboratory	Imaging	Risk score data
>1 HF hospitalization in last year - NYHA class III–IV - Intolerant of optimal dose of any GDMT HF drug - Increasing diuretic requirement - SBP <90 mmHg - Inability to perform CPET 6MWT <300 m - CRT non responder clinically - Cachexia, unintentional weight loss - KIDCO decrease >5 units	- eGFR <45 mL/min - SCr ≥160 micromoles/L - K⁺ >5.2 or <3.5 mmol/L - Hyponatremia - Hb ≤120 g/L - Persistently elevated high-BNP/NT-proBNP, e.g. NT-proBNP ≥1000 pg/mL - Abnormal liver function test - Low albumin	- LVEF ≤30% - Large area of akinesis/dyskinesis or aneurysm - Moderate/severe mitral regurgitation - RV dysfunction - Systolic PA pressure ≥50 mmHg - Moderate/severe tricuspid regurgitation - Difficult to grade aortic stenosis - IVC dilated or without respiratory variation	- MAGGIC predicted survival ≤80% at 1 year - SHFM predicted survival ≤80% at 1 year - MECKI predicted survival ≤80% at 1 year


Valeur pronostique de l'évolution du NT-proBNP

PARADIGM-HF



Zile M. J Am Coll Cardiol 2016;68:2425–36

EMPEROR-Reduced



Januzzi J. J Am Coll Cardiol 2021;78:1321–1332

IC Avancée : évaluation

- KT droit :
 - Intérêt diagnostique.

Recommendations	Class
Recommendations for the diagnosis of HF	
Right heart catheterization should be considered in patients where HF is thought to be due to congestive heart failure, restrictive cardiomyopathy, congenital heart disease, and high-output states.	IIa
Right heart catheterization may be considered in selected patients with HFpEF to confirm the diagnosis.	IIb

- Pas d'intérêt pour la sélection des patients...

- ...Mais intérêt si discussion de projet

Right heart catheterization is recommended in patients with severe HF being evaluated for heart transplantation or LVAD.	I	C
--	---	---

Contraindications

- Active infection
- Severe ventricular arrhythmia or conduction system disease
- Unstable, irreversible pulmonary hypertension (LVEDP > 30 mmHg) should be considered with subsequent re-evaluation to establish candidacy
- Cancer in collaboration with oncology specialists should occur to stratify each patient as to their risk of tumour recurrence
- Irreversible renal dysfunction (e.g. creatinine clearance <30 mL/min)
- Irreversible disease with multiple involvement
- Systemic disease with multiple involvement
- Other serious comorbidity with poor prognosis
- Pre-transplant BNP >35 ng/L (weight loss is recommended to achieve a BNP <35 ng/L)
- Current dialysis or drug abuse
- Any patient for whom social supports are deemed insufficient to achieve compliant care in the outpatient setting

HTP

1.3. Role of diagnostic right-heart catheterization
Right heart catheterization (RHC) should be performed on all candidates in preparation for listing for cardiac transplantation and annually until transplantation (Class I, Level of Evidence: C).
RHC should be performed at 3- to 6-month intervals in listed patients, especially in the presence of reversible pulmonary hypertension or worsening of heart failure symptoms (Class I, Level of Evidence: C).
A vasopressor challenge should be administered when the pulmonary artery systolic pressure is ≥ 50 mm Hg and either the transpulmonary gradient is ≥ 15 or the pulmonary vascular resistance (PVR) is > 3 Wood units while maintaining a systolic arterial blood pressure > 85 mm Hg (Class I, Level of Evidence: C).

When an acute vasopressor challenge is unsuccessful, hospitalization with continuous hemodynamic monitoring should be performed, as often the PVR will decline after 24 to 48 hours of treatment consisting of diuretics, inotropes and vasodilator agents such as inhaled nitric oxide (Class I, Level of Evidence: C).
If medical therapy fails to achieve acceptable hemodynamics, and if the left ventricle cannot be effectively unloaded with mechanical devices, including an intra-aortic balloon pump (IABP) and/or left ventricular assist device (LVAD), it is reasonable to consider that the pulmonary hypertension is irreversible. After LVAD, resolution of hemodynamics should be done after 3 to 6 months to ascertain reversibility of pulmonary hypertension (Class IIA, Level of Evidence: C).

Continuing approval without change.
If medical therapy fails to achieve acceptable hemodynamics and if the left ventricle cannot be effectively unloaded with mechanical devices, including an intra-aortic balloon pump (IABP) and/or left ventricular assist device (LVAD), it is reasonable to consider that the pulmonary hypertension is irreversible. After LVAD, resolution of hemodynamics should be done after 3 to 6 months to ascertain reversibility of pulmonary hypertension (Class IIA, Level of Evidence: C).

- PAPs >50mmHg, RVP>3UW (ou gradient TP élevé): faire test de réversibilité (TNT immédiate, si PAS « correcte », ou inotropes sur 24-48h).
- Si échec et HTAP post-capillaire pure, discuter LVAD avant la transplantation pour normalisation des pressions pulmonaires.

The 2016 International Society for Heart Lung Transplantation listing criteria for heart transplantation: A 10-year update


Cas clinique KT droit

- Homme de 64 ans, CMI diagnostiquée en 2017.
 - Projet discuté en 2018 : HTx.
 - Passage en FA, dégradation HD malgré ablation et maintien RS.
 - LVAD en bridge to transplantation fin 2018, HTx en 2021.

	07/2018	11/2018	11/2018	03/2019
AP	61/34/45	66/30/42	66/28/43	35/10/21
Pcap	37	30	30	10
OD	6	10	6	4
IC	2,5	1,8	2,3	2,8
RVP	1,8	3,8	3,2	2
Test		Levo/furo	TNT 3mg	
Post test		66/23/37, Cap 26, IC 2,1, RVP 3		


Tournants évolutifs dans l'IC

- Apparition d'arythmies (FA, TV/FV)
- Insuffisance mitrale secondaire
 - Dilatation VG, asynchronisme papillaire et tenting sous valvulaire
 - Élévation POG
- Apparition d'un syndrome cardio-rénal
 - Congestion veineuse +++
 - Bas débit, vasoconstriction artériolaire et activation SRAA
- Altération cardio-hépatique
 - Hépatopathie congestive (cholestase), ischémie hépatique (cytolysse), jusqu'à cirrhose cardiaque
- Hypertension pulmonaire post-capillaire
 - Isolée puis mixte pré et post capillaire (RVP >3)
- Défaillance VD
 - Élévation PAP et RVP, élévation PVC, altération contractile et IT




Bertrand P. Circ. 2017;135:297–314.

Quand adresser le patient ?



IC avancée : que peut-on (encore) faire ?



- TTT médicamenteux ? compliqué car svt TA basse et insuffisance rénale, souvent nécessité de ↘ TTT.
- Réadaptation pré-chirurgicale ?
- CRT ? Peu utile en cas de dysfonction VD, d'ATCD d'utilisation d'inotropes ou classe NYHA 4
- **LVAD et/ou transplantation ? Place clairement établie**
- Mitraclap ou nouvelles techniques percutanées de PEC de valvulopathies 2nd ? Place encore à déterminer
- Ultrafiltration ou dialyse péritonéale palliative ?
- Prise en charge multidisciplinaire palliative ?

Options chirurgicales

• LVAD

• Transplantation

Bridge to decision (BTD)	Use of short-term PCS (ECMO or Impella) in patients with cardiogenic shock and haemodynamic and end-organ perfusion on maximal conventional therapy for long-term PCS are evaluated (short-term bridge after resuscitation) and additional therapeutic options including long-term VAD therapy or heart transplant can be evaluated.
Bridge to candidacy (BTC)	Use of PCS (usually LVAD) to improve end-organ function and/or to make an eligible patient eligible for heart transplantation.
Bridge to transplantation (BTX)	Use of PCS (LVAD, BIVAD or TAH) to keep a patient alive who is otherwise at high risk of death before transplantation and a donor organ becomes available.
Bridge to recovery (BTR)	Use of PCS (short-term or long-term) to keep a patient alive until cardiac function recovers sufficiently to remove PCS.
Destination therapy (DT)	Long-term use of PCS (LVAD) as an alternative to transplantation in patients with end-stage HF ineligible for transplantation.

Recommendations for the treatment of patients with advanced heart failure

Recommendations	Class	Level
Patients being considered for long-term PCS must have good compliance, appropriate support for device handling and psychosocial support. ^{1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100,101,102,103,104,105,106,107,108,109,110,111,112,113,114,115,116,117,118,119,120,121,122,123,124,125,126,127,128,129,130,131,132,133,134,135,136,137,138,139,140,141,142,143,144,145,146,147,148,149,150,151,152,153,154,155,156,157,158,159,160,161,162,163,164,165,166,167,168,169,170,171,172,173,174,175,176,177,178,179,180,181,182,183,184,185,186,187,188,189,190,191,192,193,194,195,196,197,198,199,200,201,202,203,204,205,206,207,208,209,210,211,212,213,214,215,216,217,218,219,220,221,222,223,224,225,226,227,228,229,230,231,232,233,234,235,236,237,238,239,240,241,242,243,244,245,246,247,248,249,250,251,252,253,254,255,256,257,258,259,260,261,262,263,264,265,266,267,268,269,270,271,272,273,274,275,276,277,278,279,280,281,282,283,284,285,286,287,288,289,290,291,292,293,294,295,296,297,298,299,300,301,302,303,304,305,306,307,308,309,310,311,312,313,314,315,316,317,318,319,320,321,322,323,324,325,326,327,328,329,330,331,332,333,334,335,336,337,338,339,340,341,342,343,344,345,346,347,348,349,350,351,352,353,354,355,356,357,358,359,360,361,362,363,364,365,366,367,368,369,370,371,372,373,374,375,376,377,378,379,380,381,382,383,384,385,386,387,388,389,390,391,392,393,394,395,396,397,398,399,400,401,402,403,404,405,406,407,408,409,410,411,412,413,414,415,416,417,418,419,420,421,422,423,424,425,426,427,428,429,430,431,432,433,434,435,436,437,438,439,440,441,442,443,444,445,446,447,448,449,450,451,452,453,454,455,456,457,458,459,460,461,462,463,464,465,466,467,468,469,470,471,472,473,474,475,476,477,478,479,480,481,482,483,484,485,486,487,488,489,490,491,492,493,494,495,496,497,498,499,500,501,502,503,504,505,506,507,508,509,510,511,512,513,514,515,516,517,518,519,520,521,522,523,524,525,526,527,528,529,530,531,532,533,534,535,536,537,538,539,540,541,542,543,544,545,546,547,548,549,550,551,552,553,554,555,556,557,558,559,560,561,562,563,564,565,566,567,568,569,570,571,572,573,574,575,576,577,578,579,580,581,582,583,584,585,586,587,588,589,590,591,592,593,594,595,596,597,598,599,600,601,602,603,604,605,606,607,608,609,610,611,612,613,614,615,616,617,618,619,620,621,622,623,624,625,626,627,628,629,630,631,632,633,634,635,636,637,638,639,640,641,642,643,644,645,646,647,648,649,650,651,652,653,654,655,656,657,658,659,660,661,662,663,664,665,666,667,668,669,670,671,672,673,674,675,676,677,678,679,680,681,682,683,684,685,686,687,688,689,690,691,692,693,694,695,696,697,698,699,700,701,702,703,704,705,706,707,708,709,710,711,712,713,714,715,716,717,718,719,720,721,722,723,724,725,726,727,728,729,730,731,732,733,734,735,736,737,738,739,740,741,742,743,744,745,746,747,748,749,750,751,752,753,754,755,756,757,758,759,760,761,762,763,764,765,766,767,768,769,770,771,772,773,774,775,776,777,778,779,780,781,782,783,784,785,786,787,788,789,790,791,792,793,794,795,796,797,798,799,800,801,802,803,804,805,806,807,808,809,810,811,812,813,814,815,816,817,818,819,820,821,822,823,824,825,826,827,828,829,830,831,832,833,834,835,836,837,838,839,840,841,842,843,844,845,846,847,848,849,850,851,852,853,854,855,856,857,858,859,860,861,862,863,864,865,866,867,868,869,870,871,872,873,874,875,876,877,878,879,880,881,882,883,884,885,886,887,888,889,890,891,892,893,894,895,896,897,898,899,900,901,902,903,904,905,906,907,908,909,910,911,912,913,914,915,916,917,918,919,920,921,922,923,924,925,926,927,928,929,930,931,932,933,934,935,936,937,938,939,940,941,942,943,944,945,946,947,948,949,950,951,952,953,954,955,956,957,958,959,960,961,962,963,964,965,966,967,968,969,970,971,972,973,974,975,976,977,978,979,980,981,982,983,984,985,986,987,988,989,990,991,992,993,994,995,996,997,998,999,1000}		

Transplantation cardiaque

• 2 candidats pour 1 greffon

Tableau C2. Évolution des principaux indicateurs de pénurie en greffe cardiaque

Tableau C2. Évolution des principaux indicateurs de pénurie en greffe cardiaque


	2015	2016	2017	2018	2019	2020
Nouveaux inscrits pour un greffon	1,3	1,2	1,2	1,4	1,4	1,5
Receveurs en attente en liste active au 1er janvier pour un greffon*	0,5	0,6	0,5	0,4	0,6	0,7

Tableau C6. Evolution sur les trois premières années du devenir des malades inscrits pour la première fois en liste d'attente cardiaque en 2017 (N=534)

Tableau C6. Evolution sur les trois premières années du devenir des malades inscrits pour la première fois en liste d'attente cardiaque en 2017 (N= 534)

Statut sur liste d'attente (%)	à 3 mois	à 6 mois	à 12 mois	à 18 mois	à 24 mois	à 30 mois	à 36 mois
En liste inactive depuis l'inscription	1,5	0,9	0,9	0,4	0,4	0,4	0,4
En liste active	8,2	5,1	3,6	2,6	2,1	2,2	1,1
Greffé	25,8	20,2	13,3	8,8	6,2	5,2	5,4
Décédé en attente	56,2	64,4	70,2	74	76	76,8	77
Sorti de la liste d'attente pour aggravation	5,6	6,4	7,7	8,8	8,8	8,8	9,2
Sorti de la liste d'attente hors aggravation	1,5	1,5	2,1	2,2	2,4	2,4	2,6
En liste inactive depuis l'inscription et décédé ou sorti pour aggravation	0,9	1,3	1,9	3	3,9	3,9	4,1

Score cœur = Score National d'Attribution des Greffons Cardiaques (SNAGC).



Score cœur : ICAR

- Calcul de l'Index de Risque Cardiaque : ICAR
- Évaluation du risque de décès sur liste d'attente.
- Assistance de courte durée.
 - BNP ou NT-proBNP.
 - DFG par MDRD.
 - Bilirubine totale.

Sous inotropes ou assistance : valeurs AVANT implantation ou mise sous inotropes.

MISES A JOUR:

- Tous les 3 jours sous ECLS ou inotropes.
- Tous les 3 mois sinon.

Score cœur: Score Cardiaque Composite Brut



Le Score Cardiaque Composite comporte quatre composantes mutuellement exclusives (voir : ANNEXES 3.3) :

- Composante Adulte Standard
- Composante Expert Adulte (XPCA)
- Composante Pédiatrique Standard
- Composante Expert Pédiatrique (XPCP)




Tableau des composantes du Score Cardiaque Composite Brut


Composantes du score cardiaque	Nombre de points	Demande de priorité
Adulte Standard	→ 0 - 775 → 826 - 1051	⊘
Pédiatrique Standard	→ 776 - 825	⊘
Expert Adulte (XPCA)	→ 900	→ nécessaire
Expert Pédiatrique (XPCP) → Niveau1 (XPCP1) → Niveau2 (XPCP2)	→ 1102 - 1151 → 1051 - 1101	→ nécessaire

Score CCB : composantes standard

- 1 : Adulte standard ambulatoire, 0-775 points.
- 2 : Pédiatrique standard ambulatoire, (<18 ans).
- 3 : Adulte standard en choc (notamment ECLS) 826 - 1051 points.



Risque de décès en attente en fonction de la durée d'ECMO (2018)




Attention pour les patients adultes sous ECMO, les points du Score seront minorés à partir de 12 jours d'implantation (-10% par jour) et annulés dès le 16ème jour (Score=0).

Score CCB : composantes expert

Délai d'attribution du maximum de points pour la Composante Expert Adulte (XPCA) selon la situation clinique du patient

Situations cliniques pouvant faire l'objet d'une demande XPCA	Nombre de points max.	Délai d'attribution
Thrombose d'assistance circulatoire mécanique de longue durée	900	immédiat
Dysfonction d'assistance circulatoire mécanique de longue durée à l'exclusion des thromboses	900	immédiat
Orage rythmique ventriculaire non contrôlé	900	immédiat
Hémorragie chez les porteurs d'assistance circulatoire mécanique de longue durée	900	3 mois
Infection du dispositif d'assistance circulatoire mécanique de longue durée	900	3 mois
Contre-indication à l'implantation d'une assistance circulatoire mécanique de longue durée	900	3 mois
Assistance circulatoire mécanique bi ventriculaire ou cœur artificiel total	900	3 mois

Adulte expert



Attribution croissante linéaire de 0 à 900 points

Score CCB : composantes expert



Pédiatrique expert : 2 niveaux.

Composante Expert Pédiatrique - Niveaux de priorité selon la situation clinique

Situations cliniques pouvant faire l'objet d'une demande XPCP1	Situations cliniques pouvant faire l'objet d'une demande XPCP2
Assistance circulatoire de longue durée compliquée	Assistance circulatoire de longue durée non compliquée
ECMO compliquée	ECMO non compliquée
Contre-indication à la mise en place d'un Berlin Heart	Perfusion d'inotrope

Score Cardiaque Composite Pondéré



Appariement au donneur : âge, GS, S², filtrage

Age

Si le donneur est plus jeune que le receveur :

- ✓ différence d'âge ≤ 15 ans : → 100% des points
- ✓ différence d'âge ≥ 15 et ≤ 40 ans : → un pourcentage des points décroissant à partir de 100% jusqu'à 0%
- ✓ différence d'âge > 40 ans : → 0% points

Si le donneur est plus âgé que le receveur :

- ✓ différence d'âge ≤ 40 ans : → 100% des points
- ✓ différence d'âge ≥ 40 et ≤ 65 ans : → un pourcentage des points décroissant à partir de 100% jusqu'à 0%
- ✓ différence d'âge > 65 ans : → 0% points

Attribuer les greffons « jeunes » au receveurs « jeunes ».

Score Cardiaque Composite Pondéré


Age Receveur	ABO Donneur	ABO Receveur
≥ 18 ans	A	→ A, AB
	AB	→ AB
	B	→ B, AB ⁽¹⁾
	O	→ O, B
< 18 ans	A	
	AB	
	B	
	O	→ Groupe compatible

(1) Les receveurs de groupe B sont prioritaires par rapport aux receveurs de groupe AB

Surface corporelle


Type de receveur	Critères morphologiques donneur
Adulte	SCD* > 80% SCR** ou Homme ≥ 70 Kg
Pédiatrique	SCD ∈ [80% ; 300%] SCR ou Homme ≥ 70 Kg

SCD* : Surface corporelle donneur
SCR** : Surface corporelle receveur




Solution proposée pour les assistances de longue durée non compliquées


Pour les patients sous LVAD non compliqué, tous les X greffons proposés à un candidat avec moins de 750 points, octroi de 750 points au score hors appariement donneur (classement en fonction de la durée d'implantation)



Adult and Pediatric Heart Transplants Kaplan-Meier Survival (Transplants: January 1992 – June 2017)



ISHLT 2019
INTERNATIONAL SOCIETY FOR HEART AND LUNG TRANSPLANTATION



Conclusion

- ICA = pronostic sombre.
- Options thérapeutiques limitées, lourdes et projets complexes.
- Problématique de la pénurie de greffons, de la sélection des patients.
- Caractère perfectible du score cœur (CM restrictives, LVAD non compliqués...).