



## Quel objectif de pression artérielle au cours de la CEC ?


DESEBBE Olivier  
 Département d'Anesthésie, Réanimation, et Médecine périopératoire  
 Clinique de la Sauvegarde, Lyon

## Conflits d'intérêt

- MEDTRONIC
- BRAINDEX

## Définition d'une pression

$$\text{Pascal} \longleftarrow P = \frac{F}{S} \longrightarrow \text{Newton} \longrightarrow m^2$$



- Pression = Force exercée sur une surface
- agitation incessante des molécules et de leurs collisions, entre elles ou sur des obstacles.
- transfert de quantité de mouvement dans un liquide et ses effets sur des parois.
- La pression s'exerce naturellement dans toutes les directions.



## Différentes approches de la PAM

## Approche électrique: loi d'OHM

- Gradient de pression motrice  $\Delta P = DC \times RVS$

$$RVS = 8 \eta L / r^4$$

$\eta$  = viscosité sanguine vaisseaux



L = longueur du réseau

r = rayon des

La pression est générée par le débit et les résistances  
 Ex: de la CEC et l'augmentation du débit  
 Mettre vidéo


Clinical Trial > J Cardiothorac Anesth. 1990 Dec;4(6):711-4. doi: 10.1016/s0888-6296(09)90009-6.

## Histamine blockade and cardiovascular changes following heparin administration during cardiac surgery

P A Casthely <sup>1</sup>, D Yoganathan, B Kanyanis, M Saleem, T Yoganathan, C Komer, M Uribe, S Scifani, A Hudak


Affiliations + expand  
 PMID: 2131900 DOI: 10.1016/s0888-6296(09)90009-6

Vasodilatation aigue





### Histamine blockade and cardiovascular changes following heparin administration during cardiac surgery

P A Casthely<sup>1</sup>, D Yoganathan, B Karyanis, M Salem, T Yoganathan, C Komer, M Uribe, S Scifani, A Hoshik  
 Affiliations + expand  
 PMID: 2131900 DOI: 10.1016/s0888-6296(09)90009-6



### The Conundrum of Systemic Arterial Pressure Management on Cardiopulmonary Bypass


Marco Ranucci<sup>1</sup>, Mauro Cotza, and Umberto Di Dedda



$$\Delta P = \frac{Q \times 8\eta L}{\pi R^4}$$


### Heterogeneous impact of hypotension on organ perfusion and outcomes: a narrative review

Jingzhong Meng<sup>1</sup>  
 Department of Anesthesiology, Yale University School of Medicine, New Haven, CT, USA  
 email: jingzhong.meng@yale.edu



### Approche physiologique

- **Entrant:** Débit entrant = débit de CEC
- **Contenant:** Paroi des vaisseaux: élasticité, compliance artérielle
  - Élasticité
  - Impédance et inductance aortique (représente la résistance au flux pulsatile)
- **Contenu:** viscosité sanguine et plasmatique
- **Sortant:** Tonus vasomoteur distal: Résistances artérielles périphériques



Thiele A&A 2011

### Comment mesurer la pression artérielle?


Cathétérisme artériel et mesure invasive de la pression artérielle en anesthésie-réanimation chez l'adulte

Conférence d'experts - SFAR 1994

- Tête de pression en regard de la ligne axillaire moyenne...

### Quel site permet de mesurer la PAM?



- Radial/fémoral?
- PAM radiale < PAM fémorale







# Évolution de la Pression artérielle dans l'arbre vasculaire


- Gradient de pression selon Résistances et surface endothéliale



# La PAM diminue mais la pression pulsée augmente en circulation pulsatile




# Pression de perfusion capillaire De la macrocirculation vers la microcirculation




Chute de pression dans les capillaires Variable selon le tonus vasomoteur, les inotropes, les vasoconstricteurs

Pression intéressante = Pression d'entrée...



# Pression systémique moyenne (Pmsf)

Pression Systémique Moyenne pression mesurée en l'absence de flux = Pression dans les veinules postcapillaires (60% de la volémie) « obstacle aux échanges capillaires »




Joseph Anzilic, Author manuscript; available in PMC 2013 Apr 1. PMID: PMC3310203  
 Published in final edited form as: Anesth Analg. 2012 Apr; 114(4): 803-810. Published online 2012 Feb 17. doi: 10.1213/ANE.0b013e3182471e44  
 PMID: 22344243

## Determination of Vascular Waterfall Phenomenon by Bedside Measurement of Mean Systemic Filling Pressure and Critical Closing Pressure in the Intensive Care Unit

Jacinta J. Maas, M.D., Rob B. de Witte, Ph.D., Leon P. Aarts, M.D., Ph.D., Michael R. Pinsky, M.D., Dr hc, FCCM, and Job R. Jansen, Ph.D.

zone de fermeture critique qui découpe la circulation artérielle de la circulation veineuse et rend inadaptée l'utilisation de la simple PAM-PVC comme gradient de perfusion



The pressure gap of 26.8±10.7 mmHg between Pcc and Pmsf indicates that a waterfall phenomenon is likely to be present

## PAM et microcirculation

Review Article  
Monitoring of the Sublingual Microcirculation During Cardiac Surgery: Current Knowledge and Future Directions

Moritz Flick, MD<sup>1</sup>, Jacques Duranteau, MD<sup>1</sup>, Thomas W.L. Scheeren, MD<sup>1</sup>, Bernd Saugel, MD<sup>1,2</sup>

- ? Baisse du lactate
- Amélioration marqueurs de microcirculation?: SDF, ptiO2...
- Pas de marqueurs continus de la microcirculation disponibles pour la CEC

### The response of the microcirculation to cardiac surgery

Atila Kara<sup>a,b</sup>, Sakir Akin<sup>a,c</sup>, and Can Ince<sup>a</sup>

high MAP group, a higher MAP was maintained through administration of higher doses of norepinephrine and repeated bolus injections of phenylephrine. **However, there were no differences in any of the microcirculatory variables between the high and low MAP groups.**

Holmgaard Microcirculation 2018

## Pression systémique per CEC

- Résultante du tonus vasomoteur et du débit de CEC
- Pression modifiable par vasodilatateurs, vasoconstricteurs, modification du nombre de tours/min

## Influence du type de pompe sur la PAM en CEC




- principe de l'occlusion plus ou moins complète d'un tube souple par des galets rotatifs
- indépendantes de la postcharge: elles maintiennent leur débit quelle que soit la pression artérielle
- turbine rotative à haute vitesse
- sensible à la précharge et à la postcharge car non occlusive

Effet des vasoconstricteurs/vasodilatateurs sur la PAM et pas sur le débit

Effet mixte des vasoconstricteurs/vasodilatateurs sur la PAM et sur le débit

Test du tonus vasculaire sur la circulation systémique (DaO<sub>2</sub>, NIRS, SvO<sub>2</sub>, VO<sub>2</sub>...) sera plus difficile à interpréter pour la pompe centrifuge

## Quel objectif de PAM pendant la CEC?



## Hypo et hypertension artérielle per CEC



- Vasoplegic syndrome during CPB may derive from
  - the release of proinflammatory cytokines: activation coagulation, inflammation
  - anaesthetic drugs
  - active endocarditis
  - preoperative use of angiotensin-converting enzyme inhibitors and calcium channel blockers.
- Hypertension during CPB may derive from
  - an inadequate level of anaesthesia/analgesia,
  - a release of catecholamines,
  - vasoconstriction due to hypothermia and others

### 2024 EACTS/EACTAIC/EBCP Guidelines on cardiopulmonary bypass in adult cardiac surgery

Alexander Wahiba<sup>1, 2</sup>, Gudrun Kunst<sup>3, 4</sup>, Filip De Somer<sup>5</sup>, Henrik Agerup Kildahl<sup>1, 2</sup>,

## Avantages théoriques d'une PAM haute: 80-100

- Meilleure perfusion tissulaire (cerveau, rein, tube digestif) ;
- Meilleure adéquation aux besoins des malades hypertendus, âgés ou diabétiques ;
- Assurance d'un flux satisfaisant lorsque l'autorégulation d'un organe est perturbée ;
- Amélioration du flux collatéral pour les zones tissulaires à risque d'ischémie : « watershed stroke »




### Avantages d'une PAM basse

- Moins de traumatisme pour les éléments figurés ;
- Moins d'hémorragie et de retour sanguin par les artères bronchiques
- Moins de volume aspiré et lésé dans les aspirations de cardiectomie ;
- Moins de charge embolique cérébrale ;
- Meilleure protection myocardique (réduction du flux coronarien collatéral).

Pas de vasodilatateur préférentiel

"At present, the existing literature does not contain adequate information to provide recommendations on the choice of vasodilating agents."



### Association hypotension et complications postopératoires

5000 patients

**ANESTHESIOLOGY**

**EDITOR'S PERSPECTIVE**

**Intraoperative Hypotension and Acute Kidney Injury, Stroke, and Mortality during and outside Cardiopulmonary Bypass: A Retrospective Observational Cohort Study**

**What We Already Know about This Topic**

- Single-center data demonstrate that intraoperative hypotension during cardiac surgery is independently associated with stroke and acute kidney injury.
- The reproducibility of this observation and whether the timing of hypotension during cardiac surgery (within vs. outside the cardiopulmonary bypass period) modifies the association remain unclear.

**What This Article Tells Us That Is New**

- Among 4,984 patients undergoing cardiac surgery at a single tertiary care center between 2008 and 2016, 256 (5.1%) experienced the primary outcome of stroke (66, 1.3%), acute kidney injury (125, 2.5%), or mortality (109, 2.2%).
- Each 10-min of hypotension (mean arterial pressure of less than 65 mmHg) during, before, or after cardiopulmonary bypass was associated with an increased odds ratio of 1.06 (95% CI, 1.03 to 1.10; P = 0.001).
- Intraoperative hypotension, even if it occurs outside of cardiopulmonary bypass, is independently associated with stroke, acute kidney injury, or death after cardiac surgery.

The bubble chart shows the composite primary outcome rate (stroke, AKI, or mortality) for different doses of vasopressors used to treat hypotension. The x-axis is 'Vasopressor-hypotense dose' and the y-axis is 'Intraoperative hypotension total duration: MAP < 65 mmHg'. The size of the red bubbles represents the outcome rate, which increases as the hypotension duration increases.

### Les études: PAM haute ou basse?

**Circulation**

**ORIGINAL RESEARCH ARTICLE**

**High-Target Versus Low-Target Blood Pressure Management During Cardiopulmonary Bypass to Prevent Cerebral Injury in Cardiac Surgery Patients**

**A Randomized Controlled Trial**

**IMPROVEMENT OF OUTCOMES AFTER CORONARY ARTERY BYPASS**

**A randomized trial comparing intraoperative high versus low mean pressure**

Increased systemic perfusion pressure during cardiopulmonary bypass is associated with less early postoperative cognitive dysfunction and delirium

Matthias Siepe<sup>1,2</sup>, Thomas Pfeiffer<sup>1</sup>, Andreas Gieringer, Silke Zemann, Christoph Benk, Christian Schlemak, Friedhelm Beyersdorf

Department of Cardiovascular Surgery, University Medical Center Freiburg, Freiburg, Germany

**RESEARCH ARTICLE**

Higher arterial pressure during cardiopulmonary bypass may not reduce the risk of acute kidney injury


Increasing mean arterial pressure during cardiac surgery does not reduce the rate of postoperative acute kidney injury.

Matthias Siepe<sup>1,2</sup>, James C. Hebl<sup>1,2</sup>, Peter Ottavio<sup>1</sup>, Tobias E. Ammer<sup>1</sup>, Christian H. Haber<sup>1</sup>, Jens Otto Gieringer<sup>1</sup>, Frank C. Minicup<sup>1</sup>, and David A. Schemper<sup>1,2</sup>


© Author Information

**Morbidité et mortalité des patients pris en charge avec des cibles élevées ou basses de pression artérielle pendant la chirurgie cardiaque sous circulation extracorporelle : une revue systématique et une méta-analyse d'études randomisées contrôlées**

Charlotte C. McEwen, MD · Takhliliq Amir, BHSc(Hons) · Yuan Qiu, BHSc(Hons) · 2022





	n	population	Basse PAM	Haute PAM	objectif	résultats	
Azou	292	À risque d'AKI	50-60	75-85	Creat > 30%	NS	
Siepe	92	Chir coronaire	60-70	80-90	MMSE et délirium à H48	Delirium 13% vs 0%	
Gold	248	Chir coronaire	50-60	80-100	Complications neuro et cardiaques	12,9 vs 4,8% 4,0 vs 1,6% décès à 6 mois	
Vedel	197	Coronaires + valves	40-50	70-80	Volume total des lésions ischémiques cérébrales IRM POCD à J90	NS	
Miao	2022	40	Valvulaire	50-60	70-80	Lactate	3,1 versus 2,1
Kandler	2019	90	PAC Valves	40-50	> 65	AKI	NS



**Table IV. Cardiac and neurologic outcomes in the two treatment groups (intention to treat)**

	Low MAP (n = 124)		High MAP (n = 124)		Low - high MAP		95% CI for % difference
	No.	%	No.	%	No.	%	
Fatal stroke	2	1.6	0	0.0			
Hemiparesis*	2	1.6	1	0.8			
Aphasia	3	2.4	1	0.8			
Cortical blindness	1	0.8	0	0.0			
Monocular blindness	1	0.8	0	0.0			
Other focal deficit	0	0.0	1	0.8			
Total permanent neurologic complications	9	7.2	3	2.4	6	4.8	-0.5, 11.0
Fatal cardiogenic shock	1	0.8	2	1.6			
Shock	1	0.8	0	0.0			
Myocardial infarction	4	3.2	1	0.8			
Total cardiac complications	6	4.8	3	2.4	3	2.4	-2.2, 7.1
Other death, total (not attributable to cardiac or neurologic causes)	2	1.6	0	0.0	2	1.6	-0.6, 3.8
Total mortality and major cardiac and neurologic morbidity	16†	12.9	6	4.8	10	8.1	1.0, 15.1

Taux d'AC dans le groupe low MAP très élevé et grave Débit de CEC 1,9 à 2,3 l/min/m<sup>2</sup>



ORIGINAL RESEARCH ARTICLE  
**High-Target Versus Low-Target Blood Pressure Management During Cardiopulmonary Bypass to Prevent Cerebral Injury in Cardiac Surgery Patients**  
 A Randomized Controlled Trial

**Table 1. Baseline Characteristics**

	Low-Target Group (n=99)	High-Target Group (n=98)
Age, y	65.0±10.7	69.4±8.9
Male sex, n (%)	93 (93.9)	84 (85.7)
Nonwhite race, n (%)	2 (2.0)	0 (0)
Previous myocardial infarction, n (%)*	37 (37.4)	37 (37.8)
Recent myocardial infarction (past 2 wk), n (%)	30 (30.3)	25 (25.5)
Aortic valvular disease, n (%)	34 (34.3)	34 (34.7)
Angina, CCS score >1, n (%)†	64 (64.6)	47 (48.0)
Current or previous atrial fibrillation, n (%)	14 (14.1)	13 (13.3)
Hypertension, n (%)	83 (84.8)	87 (88.8)
Diabetes mellitus, type 1 or 2 (insulin treated), n (%)	10 (10.1)	10 (10.2)
Diabetes mellitus, type 2 (non-insulin treated), n (%)	14 (14.1)	14 (14.3)
Chronic lung disease, n (%)	9 (9.1)	12 (12.2)
Current smoker, n (%)	18 (18.2)	15 (15.3)
Current alcohol abuse, n (%)	7 (7.1)	7 (7.1)
BMI, kg/m <sup>2</sup> ‡	27.0 (3.8)	27.6 (4.0)
Left ventricular ejection fraction, n (%)		
>50%	54 (54.5)	50 (51.5)

ORIGINAL RESEARCH ARTICLE  
**High-Target Versus Low-Target Blood Pressure Management During Cardiopulmonary Bypass to Prevent Cerebral Injury in Cardiac Surgery Patients**  
 A Randomized Controlled Trial

	Low-Target Group (n=98)	High-Target Group (n=97)
MAP before anesthesia induction, mm Hg	92.3±15.7	96.9±13.4
MAP during bypass, mm Hg	44.7±4.7	66.8±4.9
Norepinephrine infused in the OR, µg/kg	2.65±6.01	17.43±20.14
Patients receiving norepinephrine in the OR, n (%)	35 (35.7)	90 (92.7)

ORIGINAL RESEARCH ARTICLE  
**High-Target Versus Low-Target Blood Pressure Management During Cardiopulmonary Bypass to Prevent Cerebral Injury in Cardiac Surgery Patients**  
 A Randomized Controlled Trial

Overall, diffusion-weighted imaging revealed new cerebral lesions in 52.8% of patients in the low-target group versus 55.7% in the high-target group (P=0.76). The primary outcome of volume of new cerebral lesions was

	Low-Target Group, n	High-Target Group, n	Difference (95% CI)	OR (95% CI)	P Value	
Complete cases, mean (SD) Volume AVC	89	415 (2682)	80	488 (2539)	8 (-9.78 to 994)†	0.99‡
Adverse events, n (%)						
Death	99	0 (0)	98	4 (4.1)		Inf
Creatinine, doubling of baseline value, n (%)	2 (2.0)	9 (9.4)	4.93 (1.02–48.12)	0.03		

• PAM basse et moins d'AVC, moins d'insuffisance rénale

No difference in mortality nor in the level of cognitive functioning was found according to blood pressure target during cardiac surgery long-term at 3-year follow-up.

Journal of Cardiothoracic Surgery

RESEARCH ARTICLE Open Access

**Higher arterial pressure during cardiopulmonary bypass may not reduce the risk of acute kidney injury**  
 Kristian Kandler<sup>1</sup>, Jens C. Nilsson<sup>1</sup>, Peter Olsson<sup>1</sup>, Mathias L. Jensen<sup>1</sup>, Christian H. Møller<sup>1</sup>, Jens Otto Gerneresen<sup>1</sup>, Henrik C. Aarncrup<sup>1</sup>, and Daniel A. Zweifelhofer<sup>2</sup>

38% groupe PAM 40-50 versus 46% groupe PAM > 65

90 patients

**Table 3 Postoperative data**

	CG	HPG	p value	95% CI
Delta sCr (mmol/L)	.20 (6–51)	.25 (10–49)	0.560	–18 to 18
Delta eGFR (ml/min)	–6 (–15–4)	–3 (–13–5)	0.522	–7 to 5
<b>AKI</b>	<b>16 (50%)</b>	<b>19 (60%)</b>	<b>0.447</b>	
Dialysis	4 (10%)	3 (7%)	0.565	
Re-operation	4 (10%)	1 (3%)	0.165	
Stroke	3 (8%)	2 (5%)	0.687	
4 months follow-up				
Change in GFR (ml/min)	–9 ± 12	–5 ± 16	0.288	–13 to 4
> 10% decrease in GFR*	11 (44%)	9 (39%)	0.732	
30-day mortality	3 (7%)	3 (7%)	1.000	
6-month mortality	7 (16%)	8 (18%)	0.763	

Percentages are given as total within group. Continuous data are presented as mean ± SD or median (interquartile min – max).  
 AKI Acute kidney injury, CI confidence interval, AP Arterial pressure, eGFR Estimated glomerular filtration rate (calculated using the Cockcroft-Gault formula), GFR Glomerular filtration rate (estimated by Cr-EDTA clearance), sCr Serum creatinine.  
 \*Percentages of total of patients at follow-up.

# Prévention de l'AKI par la PAM?

SYSTEMATIC REVIEW ARTICLE  
 Society of Cardiovascular Anesthesiologists Clinical Practice Update for Management of Acute Kidney Injury Associated With Cardiac Surgery

Yu Peng, MD, PhD,<sup>1</sup> David R. Mottus, MBBS,<sup>2</sup> Bruce A. Bollen, MD,<sup>3</sup> Frederic T. Billings, RC MD,<sup>4</sup> Alexander Zamboni, MD,<sup>5</sup> William M. Popovich, MD,<sup>6</sup> Kenneth A. Fox, MD,<sup>7</sup> William A. Gattuso, MD,<sup>8</sup> Shuangling Zhou, MD,<sup>9</sup> Maria A. Gendy, MD,<sup>10</sup> Rajaraj S. MD, PhD,<sup>11</sup> Meera Bhutta, MD,<sup>12</sup> Benjamin M. Schmitt, MD,<sup>13</sup> Andrew D. Shaw, MD,<sup>14</sup> FICCA, FRCR,<sup>15</sup> FRCR,<sup>16</sup> and Hong Lu, MD,<sup>17</sup>

**Table 1. Study Characteristics**

Study	Region	Procedure	N	Interventional and control treatments, and main findings on CS-AKI	AKI definition	Journal (IF2020)
Azu et al 2014 <sup>18</sup>	France	CABG, valve, or aorta, with CPB	292	A high level of MAP (75–85 mm Hg) versus control (MAP 70–80 mm Hg) during normothermic CPB did not reduce the risk of AKI.	30% rise in sCr	Perfusion (1.97)
Kandler et al 2019 <sup>19</sup>	Denmark	CABG + valve, with CPB	90	Arterial pressure <60 mmHg versus control (MAP 47 mm Hg) during CPB did not reduce the incidence of AKI or chronic kidney injury at postoperative 4 mo.	RIFLE	J Cardiothorac Sur (1.64)
Vedell et al 2018 <sup>20</sup>	Denmark	CABG and/or valve, with CPB	197	A higher MAP (70–80 mm Hg) versus control (MAP 40–50 mm Hg) increased the number of patients with postoperative doubling of sCr.	sCr > 2 times of baseline	Circulation (26.7)

# Métaanalyse 2022

Outcome (number of studies)	High MAP target, subtotal N	Low MAP target, subtotal N	Relative risk	95% CI	P value	I <sup>2</sup> (%)
<b>Dichotomous variables</b>						
Hospital mortality (4)	8/111	6/316	1.1	0.4 to 3.3	0.84	0
30-day mortality (4)	12/334	6/337	1.6	0.6 to 4.36	0.33	0
6-month mortality (3)	15/310	19/309	0.8	0.4 to 1.6	0.55	0
Myocardial infarction (3)	7/367	8/367	0.9	0.3 to 3.2	0.89	18
Delirium (2)	10/142	13/147	0.5	0.03 to 8.6	0.61	75
Cognitive decline (2)	39/190	35/204	1.2	0.7 to 2.1	0.46	35
Stroke (5)	11/261	13/266	1.0	0.2 to 4.9	0.95	65
Hemodialysis (5)	11/368	11/372	1.0	0.5 to 2.3	0.98	0
RIFLE "risk" criteria (4)	120/297	98/303	1.3	0.9 to 1.8	0.25	41
Number transfused (3)	85/227	60/229	1.4	1.1 to 1.9	0.004	0
<b>Outcome (number of studies) High MAP target, mean (SD) Low MAP target, mean (SD) Weighted mean difference 95% CI P value I<sup>2</sup> (%)</b>						
<b>Continuous variables</b>						
ICU LOS in days (4)	2.4 (4.7)	2.6 (5.3)	0.2	–0.4 to 0.9	0.55	29
Hospital LOS in days (4)	10.7 (7.8)	11.1 (9.4)	1.1	0.3 to 1.8	0.006	13
PRBCs transfused/patient (3) 2.1 (2.7)	1.9 (2.9)	0.1	–0.1 to 0.3	0.35	0	

CI = confidence interval; ICU = intensive care unit; MAP = mean arterial pressure; LOS = length of stay; PRBC = packed red blood cell; RIFLE = Risk, Injury, Failure, Loss, and End-stage Kidney classification.

## High versus low blood pressure targets for cardiac surgery while on cardiopulmonary bypass

Yuki Kotani<sup>1</sup>, Yuki Kataoka<sup>2,3,4,5</sup>, Junichi Izawa<sup>6,7</sup>, Shoko Fujieka<sup>8</sup>, Takuo Yoshida<sup>9,10</sup>, Junji Kumasawa<sup>11,12</sup>, Joey Sw Kivong<sup>13</sup>

Affiliations + expand  
PMID: 36448514 PMCID: PMC9709767 (available on 2023-11-30)  
DOI: 10.1002/14651858.CD013494.pub2

### Conclusions des auteurs

Une cible de pression artérielle élevée pourrait n'entraîner que peu ou pas de différence dans les critères de jugement relatifs aux patients, y compris les lésions rénales aiguës et la mortalité. Compte tenu de largeur des intervalles de confiance, d'autres études sont nécessaires pour confirmer l'efficacité d'une cible de pression artérielle plus élevée chez les personnes qui subissent une chirurgie cardiaque avec pontage cardio-pulmonaire.

## Reco EACTS 2024 sur la PAM per CEC

Recommendation Table 29 Recommendations for kidney protection

Recommendations	Class <sup>a</sup>	Level <sup>b</sup>	Ref <sup>c</sup>
Routinely targeting a high MAP using vasoconstrictors is not recommended during CPB to reduce AKI.	III	A	377,378,380,381

## Reco EACTS 2024


Recommendation Table 40 Recommendations for control of mean arterial blood pressure during cardiopulmonary bypass

Recommendations	Class <sup>a</sup>	Level <sup>b</sup>	Ref <sup>c</sup>
It is recommended that the MAP be maintained between 50 and 80 mmHg with vasoconstrictors and vasodilators if required, having ensured that the depth of anaesthesia and pump flow rate are sufficient.	I	A	381,511
The use of vasopressors to increase the MAP to values above 80 mmHg during CPB is not recommended.	III	B	381,510,517
Targeting the MAP during CPB within the limits of individualized cerebral autoregulation data, measured under normocapnic conditions before CPB, should be considered whenever the technical and human skills are available.	IIa	A	222,519,520
It is recommended that vasoplegic syndrome during CPB be treated with $\alpha$ 1-adrenergic agonists and/or vasopressin.	I	C	521,523
In refractory vasoplegic syndrome, alternative drugs (methylene blue or terlipressin) should be considered, alone or in combination.	IIa	B	522,523
Hydroxocobalamin or angiotensin II may be considered to treat vasoplegic syndrome during CPB.	IIb	C	524–527

## Titrer la PAM pendant la CEC?

= individualiser pour chaque patient le niveau de PAM selon ses besoins


## Détermination physiologique de la PAM idéale



## Optimisation de la PAM? Autorégulation de l'organe

Pas de variation du DSR et du DFG pour des variations de pression artérielle moyenne entre 80 et 180 mmHg

Cerveau  
Rein  
Myocarde (avant/après clampage)  
Mésentère?



## Titration selon la valeur initiale de PAM?

**Improvement of Outcomes after Coronary Artery Bypass II: A Randomized Trial Comparing Intraoperative High Versus Customized Mean Arterial Pressure**  
 Mary E. Charlton, M.D., Janey C. Peterson, E.D., R.N., Karl H. Kröger, M.D.

Adaptation PAM per CEC selon valeur basale PAM

- Sédation, pose KT artériel, attente 4 min
- the next 6 mean blood pressures downloaded to the computer from the operating room monitor (at 10 sec intervals) were recorded and an average MAP was taken.

**Improvement of Outcomes after Coronary Artery Bypass II: A Randomized Trial Comparing Intraoperative High Versus Customized Mean Arterial Pressure**  
 Mary E. Charlton, M.D., Janey C. Peterson, E.D., R.N., Karl H. Kröger, M.D.

- 412 patients
- Chir coronaire programmée
- 80 +/- 20 mmHg,
- versus
- max 90 (57-90), 88 moy
- Complications neuro et cardiaques/cognitives: NS 16,5 vs 14,5%

- Among those whose average MAP was more than 20 mmHg below their autoregulatory range, the cardiac and neurologic complication rate was double that of patients whose MAP was not that low (15.9% vs. 7.3%).
- 61% adhérence gp PAM préop
- 85% adhérence gp Pam 80
- management at higher MAP on CPB dramatically reduces stroke occurrence in patients with severe aortic atheromatous disease (grade IV and V)

**Improvement of Outcomes after Coronary Artery Bypass II: A Randomized Trial Comparing Intraoperative High Versus Customized Mean Arterial Pressure**  
 Mary E. Charlton, M.D., Janey C. Peterson, E.D., R.N., Karl H. Kröger, M.D.

## Même PAM dans les 2 groupes...

B. Intraoperative Management	High MAP	Custom MAP	P
<b>Pre-bypass</b>			
Pre-bypass MAP (mmHg)	86 ± 8	85 ± 9	
Pre-bypass cardiac output (L/min)	4 ± 1	4 ± 1	
Bypass			
Times (min)			
Bypass	74 ± 24	77 ± 22	ns
Cross-clamp	40 ± 15	40.4 ± 14	ns
Pump flows (L/min/m <sup>2</sup> ) <sup>#</sup>			
Bypass on—warming (± #)	2.0 ± 0.3	2.0 ± 0.3	ns
Warming—cross-clamp off (± #)	2.1 ± 0.3	2.2 ± 0.3	ns
Cross-clamp off—bypass off (± #)	2.4 ± 0.2	2.34 ± 0.3	ns

## observationnel

### Difference between pre-operative and cardiopulmonary bypass mean arterial pressure is independently associated with early cardiac surgery-associated acute kidney injury

Delta MAP > 26 mmHg

Hussein D Kanji, Costas J Schulze, Marilou Hervas-Malo, Peter Wang, David B Ross, Mohamad Zibdawi and Sean M Bagshaw

*Journal of Cardiothoracic Surgery* 2010 5:71  
<https://doi.org/10.1186/1749-8090-5-71> © Kanji et al; licensee BioMed Central Ltd. 2010  
 Received: 7 May 2010 Accepted: 8 September 2010 Published: 8 September 2010

Nonpulsatile pump flow rates were kept at 2.4 L/min/m<sup>2</sup>

Mesurer en site fémoral si petit gabarit, procédure longue

Position de la tête de pression...

### Difference between pre-operative and cardiopulmonary bypass mean arterial pressure is independently associated with early cardiac surgery-associated acute kidney injury

**Table 4 Multi-variable adjusted logistic regression model<sup>#</sup> of association between delta MAP and CSA-AKI**

Parameter	Odds Ratio	95% CI	P-value
Male sex	0.7	0.3-1.7	0.49
Age ≥75 years (present)	2.1	0.9-4.9	0.08
BMI ≥25 kg/m <sup>2</sup> (present)	4.2	1.6-11.2	0.0039
Delta MAP ≥26 mmHg (present)	2.8	1.3-6.1	0.009
Flow ≥54 per mL/kg/min (present)	0.3	0.1-0.7	0.004
Side-biting clamp (present)	3.0	1.3-7.1	0.012


Abbreviations: BMI = Body Mass Index; MAP = mean arterial pressure; CPB = cardiopulmonary bypass  
 Model characteristics: C-statistic = 0.788



**Automated Ambulatory Blood Pressure Measurements and Intraoperative Hypotension in Patients Having Noncardiac Surgery with General Anesthesia**

A Prospective Observational Study

Bernd Saugel, M.D., Philip C. Reese, M.D., Daniel I. Sessler, M.D., Christian Burfeindt, Julia Y. Nicklas, M.D., Hans O. Painschmidt, Ph.D., Daniel A. Reuter, M.D., Stefan Südfeld, M.D.  
ANESTHESIOLOGY 2019; 131:74-83




Nighttime, lowest value

**Automated Ambulatory Blood Pressure Measurements and Intraoperative Hypotension in Patients Having Noncardiac Surgery with General Anesthesia**




A Prospective Observational Study

Bernd Saugel, M.D., Philip C. Reese, M.D., Daniel I. Sessler, M.D., Christian Burfeindt, Julia Y. Nicklas, M.D., Hans O. Painschmidt, Ph.D., Daniel A. Reuter, M.D., Stefan Südfeld, M.D.  
ANESTHESIOLOGY 2019; 131:74-83




**Fig. 6.** Difference between lowest nighttime mean arterial pressure (MAP) and a MAP of 65 mmHg. Histogram showing the frequency (in y axis; n total = 370) of the difference between the lowest nighttime MAP and a MAP of 65 mmHg. The median difference (25th to 75th percentile range) was +5 (-1 to +13) mmHg. The dotted vertical line represents the median difference.

Ajuster la PAM selon la circulation cérébrale?




Enregistrement continu  
Analyse sur logiciel dédié de données PAM/DSC  
Corrélation de Pearson temps (5 min/30 min): **Mx**



### Optimal blood pressure during cardiopulmonary bypass defined by cerebral autoregulation monitoring



Daijiro Hori, MD,<sup>a</sup> Yohei Nomura, MD,<sup>a</sup> Masahiro Ono, MD,<sup>b</sup> Brijen Joshi, MD,<sup>c</sup> Kaushik Mandal, MD,<sup>d</sup> Duke Cameron, MD,<sup>e</sup> Masha Kocherginsky, PhD,<sup>d</sup> and Charles W. Hogue, MD<sup>e</sup>




**FIGURE 2.** Representative graph of cerebral autoregulation monitoring by Mx during CPB. The top graph represents the time-series of ABP, and the bottom bar-graph represents the percentage of the time of the recording spent at 5 mmHg bin. Optimal MAP (ABP) for the left and right side of the brain was defined as that MAP with the lowest Mx. LLA and ULA were defined as the MAP at which Mx reached 0.4. In this example, the optimal MAP is 85 mm Hg (black arrow) and the LLA and ULA are 70 mm Hg and 95 mm Hg, respectively (black-outlined arrows). ABP, Arterial blood pressure.

### Variabilité individuelle de la LLA

March 2012 • Volume 114 • Number 3

### Predicting the Limits of Cerebral Autoregulation During Cardiopulmonary Bypass


Brijen Joshi, MD,<sup>a</sup> Masahiro Ono, MD,<sup>b</sup> Charles Brown, MD,<sup>c</sup> Kenneth Brady, MD,<sup>d</sup> R. Blaine Easley, MD,<sup>e</sup> Gayane Yenokyan, PhD,<sup>f</sup> Rebecca F. Gottesman, MD, PhD,<sup>g</sup> and Charles W. Hogue, MD<sup>h</sup>



**Figure 3.** Number of subjects versus the mean arterial blood pressure at the lower limit of cerebral blood flow autoregulation during cardiopulmonary bypass based on the transcranial Doppler-determined mean velocity ratio.

www.anesthesia-analgesia.org

### Détermination du LLA avec l'oxymétrie cérébrale



**Figure 5.** Distributions of LLAs—cotrending algorithm and TCD reference. A, Bland-Altman plot of the data displaying mean bias = 0.22 mmHg, an upper LOA = 10.79 mmHg (95% CI, 10.09–11.49) and a lower LOA = -10.35 mmHg (95% CI, -9.65 to -11.05). B, Scatter plot of individual LLAs over time over all cases (bubble sizes proportional to number of collocated data points). The dashed line in the middle represents the unity line, which indicates a perfect agreement between the method and the reference. CI indicates confidence intervals; LLA, lower limit of autoregulation; LOA, limit of agreement.

### Les études observationnelles


#### Blood Pressure Excursions Below the Cerebral Autoregulation Threshold During Cardiac Surgery Are Associated With Acute Kidney Injury

Masahiro Ono, MD, PhD,<sup>1</sup> George J. Amatoakis, MD,<sup>1</sup> Derek M. Fine, MD,<sup>1</sup> Kenneth Brady,

Published in final edited form as: *J Clin Anesth*. 2014 January; 14(1): 1-6. doi:10.1016/j.jclin.2013.07.009.

Duration and magnitude of blood pressure below cerebral autoregulation threshold during cardiopulmonary bypass is associated with major morbidity and operative mortality


Masahiro Ono, MD, PhD,<sup>1</sup> Kenneth Brady, MD,<sup>1</sup> R. Blaine Easley, MD,<sup>1</sup> Charles Brown,



Variable	No MIMOM (n = 354)	MIMOM (n = 83)	P value
Average MAP during CPB (mm Hg)	74 ± 8 (73–75)	75 ± 9 (72–76)	.203
Average rScO <sub>2</sub>	54 ± 11 (52–55)	55 ± 7 (53–56)	.388
Average COX	0.27 ± 0.18 (0.25–0.29)	0.26 ± 0.17 (0.21–0.29)	.749
LLA (mm Hg)	69 ± 14 (67–70)	71 ± 12 (67–72)	.136
AUC <sub>MAP-LLA</sub> (mm Hg × min/h)	2.4 (1.1–5.7)	6.5 (2.1–15.4)	.017

### Arterial pressure above the upper cerebral autoregulation limit during cardiopulmonary bypass is associated with postoperative delirium

D. Hori<sup>1</sup>, C. Brown<sup>2</sup>, M. Ono<sup>3</sup>, T. Rappold<sup>4</sup>, F. Sieber<sup>2</sup>, A. Gottschalk<sup>2</sup>, K. J. Neufeld<sup>3</sup>, R. Gottesman<sup>1</sup>, H. Adachi<sup>5</sup> and C. W. Hogue<sup>2\*</sup>



## 2 études prospectives

### JAMA Surgery | Original Investigation Effect of Targeting Mean Arterial Pressure During Cardiopulmonary Bypass by Monitoring Cerebral Autoregulation on Postsurgical Delirium Among Older Patients A Nested Randomized Clinical Trial


Charles W. Hogue<sup>1</sup>, MD, MPH, Kevin J. Neufeld<sup>2</sup>, MD, MPH, Bing Tian<sup>3</sup>, MS, Julia Probert<sup>4</sup>, BA, Andrew Laffan<sup>5</sup>, BA, Laura Mac<sup>6</sup>, MHS, PA-C, DiploMed, MD, Yoshi Nomura<sup>7</sup>, MD, Kazuhiko Maruyama<sup>8</sup>, MD, Kenji Sudo<sup>9</sup>, MD, Charles H. Hogue<sup>10</sup>, MD, and the Cerebral Autoregulation Study Group

Randomized Controlled Trial | *Semin Thorac Cardiovasc Surg*. 2021; Summer33(2):429-438. doi: 10.1053/j.semtovs.2020.09.032. Epub 2020 Nov 10.

### Personalized Blood Pressure Management During Cardiac Surgery With Cerebral Autoregulation Monitoring: A Randomized Trial

Charles W. Hogue<sup>1</sup>, Charles H. Brown 4th<sup>2</sup>, Dajiro Hori<sup>3</sup>, Masa Ono<sup>4</sup>, Yoshi Nomura<sup>3</sup>, Lauren C. Balmert<sup>5</sup>, Nina Sclarowicz<sup>6</sup>, Jordan Grafman<sup>7</sup>, Kenneth Brady<sup>8</sup>, Cerebral Autoregulation Study Group

Affiliations + expand  
PMID: 33186735 DOI: 10.1053/j.semtovs.2020.09.032



The lower limit of autoregulation was determined by the senior author (C.W.H.) before CPB based on the highest MAP where Mx increased from less than 0.4 to 0.4 or greater. When Mx did not cross 0.4 clearly, the lower limit of autoregulation was defined as the blood pressure with the lowest Mx (the MAP with the best autoregulation)

Figure 2. Delirium Incidence by Randomization Group




Table 2. Characteristics of Management During Cardiopulmonary Bypass for Patients Randomized to Standard Care vs Autoregulation-Targeted Management of Mean Arterial Pressure

Characteristic of Management	Standard Care (n = 94)	Autoregulation-Targeted (n = 105)	P Value
Phenylephrine, median (IQR), mg	1.2 (0.3-2.3)	1.8 (0.5-3.6)	.02
Vasopressin administration, No. (%)	6 (6.4)	9 (8.6)	.56
Cardiopulmonary bypass flow, mean (SD), L/min	4.4 (0.6)	4.4 (0.6)	.92
Isoflurane, mean (SD), %	0.76 (0.27)	0.77 (0.31)	.71
Arterial pressure during cardiopulmonary bypass, mean (SD), mm Hg	71.3 (7.6)	73.9 (6.7)	.01
Arterial pressure at the lower limit of autoregulation, mean (SD), mm Hg	68.7 (11.3)	66.0 (10.9)	.10
Product of the duration of time and mean arterial pressure below the lower limit of autoregulation, median (IQR), mm Hg × hr	9.5 (3.7-19.5)	5.3 (2.0-13.4)	.002


Mean arterial pressure during cardiopulmonary bypass was managed to standard care or autoregulation-targeted goals. Delirium incidence randomization group is shown.

### Seminars in Thoracic and Cardiovascular Surgery Volume 33, Issue 2, Summer 2021, Pages 429-438

### Personalized Blood Pressure Management During Cardiac Surgery With Cerebral Autoregulation Monitoring: A Randomized Trial

Charles W. Hogue MD, MPH, Kevin J. Neufeld MD, MPH, Bing Tian MS, Julia Probert BA, Andrew Laffan BA, Laura Mac MHS, PA-C, DiploMed, MD, Yoshi Nomura MD, Kazuhiko Maruyama MD, Kenji Sudo MD, Charles H. Hogue MD, and the Cerebral Autoregulation Study Group

AVC, troubles cognitifs



a reduction in the frequency of delirium and better performance on tests of memory 4-6 weeks after surgery

Limites de l'approche de la PAM titrée sur la circulation cérébrale

### Hemodilution Combined With Hypercapnia Impairs Cerebral Autoregulation During Normothermic Cardiopulmonary Bypass

Ervin E. Sveredija, EXP<sup>1</sup>, Nussajka P.A. Vranckan, B.A.Sc.<sup>2</sup>, Antonio P. Simons, EXP, PhD,<sup>3</sup> Erik D. Gommer, MSc, PhD,<sup>4</sup> John H. Najjar, MD, PhD,<sup>5</sup> Jose G. Masson, MD, PhD,<sup>6</sup> and Patrick W. Veenendaal, EXP, PhD<sup>7</sup>






Fig 1. Cerebral autoregulation as indicated by the autoregulation index (ARI) at baseline, at 3 levels of p<sub>a</sub>CO<sub>2</sub>, and at 2 levels of hemocrit during cardiopulmonary bypass.

## Effet de la PaCO2 et de la PaO2 sur le DSC



## Altération de l'AC et réchauffement...



## Limites de l'optimisation de la PAM sur le doppler TC

- Technique à apprendre: ACM?, position
- Matériel spécifique: connecter en continu le signal de PAM et le doppler/NIRS
- LLA évolutive au cours du temps et des conditions: température, pulsatilité, PaCO2, Hte,

## Études en cours

### Trial to Compare Different Strategies of Mean Arterial Pressure Management During Cardiopulmonary Bypass

Integrated University Hospital Trust of Verona  
Status: Enrolling  
Randomized Controlled Trial  
Am Heart J. 2023 Jul 28;170:20. doi: 10.1016/j.ahj.2023.03.005.  
Epub 2023 Mar 18.

### Perioperative individualized hemodynamic optimization according to baseline mean arterial pressure in cardiac surgery patients: Rationale and design of the OPTIPAM randomized trial

Richard Descamps<sup>1</sup>, Julien Ancelet<sup>2</sup>, Emmanuel Becher<sup>3</sup>, Julien Sougbe<sup>4</sup>, Hélène Charbonneau<sup>5</sup>, Martin Charvin<sup>6</sup>, Benjamin Genty<sup>7</sup>, Olivier Desobry<sup>8</sup>, Jean-Louis Leger<sup>9</sup>, Cyril Fagot<sup>10</sup>, François Labadie<sup>11</sup>, Jean-Louis Leger<sup>12</sup>, Yacine Mahjoub<sup>13</sup>, Paul Michel<sup>14</sup>, Serge Muller<sup>15</sup>, Pierre-Henri Pagan<sup>16</sup>, Abdoumalik Diallo<sup>17</sup>, Jean-François Chénier<sup>18</sup>, Alexandre Clavier<sup>19</sup>, Sophie Provencher<sup>20</sup>, Bernard Bozer<sup>21</sup>, Jean-Jacques Pellerin<sup>22</sup>, Marc-Olivier Fischer<sup>23</sup>; OPTIPAM Investigators

Trials. 2024 Mar 15;25(1):60. doi: 10.1186/s13063-024-07792-0

### Mean Arterial Pressure (MAP) Trial: study protocol for a multicentre, randomized, controlled trial to compare three different strategies of mean arterial pressure management during cardiopulmonary bypass

Alexandra Franca<sup>1,2\*</sup>, Gina Mazzeo<sup>1</sup>, Antonella Galone<sup>1</sup>, Daniele Livardi<sup>1</sup>, Lúcio San Biagio<sup>1</sup>, Giovanni Battista Luciani<sup>1</sup>, Francesco Onorati<sup>1</sup>

\* Author information • Article notes • Copyright and License Information  
PMCID: PMC10941373 PMID: 38591502

**Trial design**  
500 participants in 3 patient groups

Standard MAP	High MAP
<b>Other group:</b> Second Comparator group MAP comparable to the patient's pre-operative MAP. This trial will be conducted by performing 2 blood pressure measurement in three different treatment of the trial before surgery (SB, BM, H) 2 hrs, and 4 hrs, and will be conducted during the standard form on 2000cc AP + 3.33 L (Lysine) AP + 80cc H2O. The preoperative MAP will be defined as the target during CPB, within a range of a 10 mmHg increase.	<b>Other group:</b> First Comparator group: MAP values between 70-80 mmHg. <b>Intervention:</b> Other Management of mean arterial pressure during cardiopulmonary bypass in cardiac surgery interventions. The MAP will be maintained in the range of values of each study arm.


## Notre expérience ...

### Assessment of Cerebral Autoregulation Patterns with Near-infrared Spectroscopy during Pharmacological-induced Pressure Changes

Annelies T. Moerman, M.D., Ph.D., Valérie M. Vanbiervliet, M.D., Astrid Van Wesselmaer, M.D., Stefan M. Bouchez, M.D., Patrick F. Wouters, M.D., Ph.D., Stefan G. De Hert, M.D., Ph.D.



## Limites de la NIRS




**Impact of 2 Distinct Levels of Mean Arterial Pressure on Near-Infrared Spectroscopy During Cardiac Surgery: Secondary Outcome From a Randomized Clinical Trial**

Christoph Henningsen<sup>1,2,3\*</sup>, Jochen S. Hinkel<sup>4,5</sup>, Christof Gellera<sup>6,7,8</sup>, Hans-Joachim Schmack<sup>9,10,11</sup>, Jan C. Nilsson<sup>12,13,14</sup>, and Henrik B. Ross<sup>15,16,17,18,19,20,21</sup>

### Cutaneous Vasoconstriction Affects Near-infrared Spectroscopy Determined Cerebral Oxygen Saturation during Administration of Norepinephrine

Henrik Sorensen, M.Sc.,\* Niels H. Secher, M.D., D.M.Sc.,† Christoph Siebenmann, M.Sc.,‡ Henning B. Nielsen, M.D., D.M.Sc.,§ Matthias Kohl-Bareis, Ph.D.,|| Carsten Lundby, Ph.D.,# Peter Rasmussen, Ph.D.\*\*



Mort encéphalique: pas de perfusion sur la carotide interne  
Clampage aortique:

Peroperative Medicine | February 2024

**Effect of Extracerebral Contamination on Near-infrared Spectroscopy as Revealed during Organ Donation: A Prospective Observational Study in Brain-dead Organ Donors**

Marie Sophie, M.D., Ph.D., M.A.B.A.; Ludovic Legeay, M.D.; Etienne Wehrli-Schneiter, M.D., Ph.D.; Stefan Oberholzer, M.D., Ph.D.; Hans-Claudio Kuehni, Ph.D.; Johannes Thurnher, M.D., Ph.D.

Anesthesiology February 2024, Vol 140, 220-229.

**Marqueur de perfusion tissulaire**  
**Manque de spécificité de monitoring du tissu cérébral**

### Intégrer le débit de CEC dans la réflexion sur la PAM

- PAM = DC \* RVS\* k

Cerebral Oximetry and Mean Arterial Pressure: Not a Straight Relationship, the Flow Between?

Desebbe O, et al. Anesth Analg 2019. PMID 31743211

#### Circulation

#### ORIGINAL RESEARCH ARTICLE

#### High-Target Versus Low-Target Blood Pressure Management During Cardiopulmonary Bypass to Prevent Cerebral Injury in Cardiac Surgery Patients: A Randomized Controlled Trial

In 1928, Jarisch is quoted as saying, "It is a source of regret that the measurement of flow [i.e., SV] is so much more difficult than the measurement of pressure. This has led to an undue interest in the blood pressure manometer. Most organs, however, require flow rather than pressure."

Table 2. Intraoperative Management

	Low-Target Group (n=98)	High-Target Group (n=97)
Hematocrit, before start of surgery, %	40.3±5.9	40.6±4.7
MAP before anesthesia induction, mm Hg	92.3±15.7	96.9±13.4
MAP during bypass, mm Hg	44.7±4.7	66.8±4.9
MAP below target during bypass, n (%) <sup>a</sup>	2 (2.0)	18 (18.5)
MAP above target during bypass, n (%) <sup>b</sup>	5 (5.1)	0 (0)
Blood flow rate during bypass, L·min <sup>-1</sup> ·m <sup>-2</sup>	2.69±0.1	2.69±0.1
Hematocrit, mean level during bypass, %	31.5±3.8	33.1±4.2
Nadir hematocrit sampling value during bypass, %	28.7±3.7	29.2±4.0
Surgery time, min	184.9±50.8	194.3±66.6
Bypass time, min	94.0±33.0	105.6±77.4
Cross-clamp time, min <sup>c</sup>	63.3±26.9	64.8±32.6
Peak lactate level during surgery, mmol	2.25±0.83	2.16±0.82
Norepinephrine infused in the OR, µg/kg	2.65±6.01	17.43±20.14
Patients receiving norepinephrine in the OR, n (%)	35 (35.7)	90 (92.7)

### Improvement of Outcomes after Coronary Artery Bypass II: A Randomized Trial Comparing Intraoperative High Versus Customized Mean Arterial Pressure

Mary E. Charlson, M.D.,\* Janey C. Peterson, E.D., B.N.,\* Karl H. Kiloeger, M.D.,†

### Débits de CEC bas...

B. Intraoperative Management Pre-bypass	High MAP	Custom MAP	P
Pre-bypass MAP (mmHg)	86 ± 8	85 ± 9	
Pre-bypass cardiac output (L/min)	4 ± 1	4 ± 1	
Bypass Times (min)	74 ± 24	77 ± 22	ns
Bypass Cross-clamp	40 ± 15	40.4 ± 14	ns
Pump flows (L/min/m <sup>2</sup> ) <sup>a</sup>			
Bypass on—warming (± #)	2.0 ± 0.3	2.0 ± 0.3	ns
Warming—cross-clamp off (± #)	2.1 ± 0.3	2.2 ± 0.3	ns
Cross-clamp off—bypass off (± #)	2.4 ± 0.2	2.34 ± 0.3	ns

	n	population	Basse PAM	Haute PAM	Débit (l/min/m <sup>2</sup> )	particularités	
Azau	2014	300	À risque d'AKI	50-60	75-85	2,4 pour SvO <sub>2</sub> > 70% 2,57 / 2,58	Circuit clot, miniCEC
Siepe	2011	92	Chir coronaire	60-70	80-90	2,6	Troubles cognitifs
Charlson	2007	412	Chir coronaire programmée	80	+/- 20 mmHg, max 90 (57-90), 88 moy	2 à 2,4	61% adhérence gp PAM préop 85% adhérence gp Pam 80 management at higher MAP on CPB dramatically reduces stroke occurrence in patients with severe aortic atherosclerotic disease (grade IV and V)
Gold	1995	248	Chir coronaire	50-60	80-100	1,9 à 2,3	Niveau de PAM respectés
Vedel	2018	197	Coronaires + valves	40-50	70-80	2,7	

le débit de pompe est primordial

# Le débit de CEC avant la pression ?...

Observational Study > J Cardiothorac Vasc Anesth, 32 (2), 684-690 Apr 2018

## Disturbances in Oxygen Balance During Cardiopulmonary Bypass: A Risk Factor for Postoperative Delirium

Nina Smulter<sup>1</sup>, Helena Claesson Lingehall<sup>2</sup>, Yngve Gustafson<sup>3</sup>, Birgitta Olofsson<sup>4</sup>, Karl Gunnar Engström<sup>5</sup>, Micael Appelblad<sup>6</sup>, Staffan Svenmarker<sup>6</sup>

Ranucci et al Perioperative Management

## Goal-directed perfusion to reduce acute kidney injury: A randomized trial

Marco Ranucci, MD, FESC,<sup>a</sup> Ian Johnson, CCP,<sup>b,c</sup> Timothy Willcox, CCP,<sup>b,c</sup> Robert A. Baker, PhD, CCP,<sup>d</sup>




FIGURE 2. Oxygen delivery ( $DO_2$ ) values in the goal-directed perfusion (blue; A) and control (red; B) arms during cardiopulmonary bypass (CPB). Boxes represent interquartile range, lines in the boxes represent the median, whiskers are 95% confidence intervals, and dots are outliers. The green line represents the critical  $DO_2$  value of  $280 \text{ mL} \cdot \text{min}^{-1} \cdot \text{m}^{-2}$ . Data are restricted to the first 120 minutes of CPB.

Etude multicentrique, Valeurs hautes de DaO2 dans les 2 bras.... Et pourtant moins d'insuffisance rénale Aigue dans le groupe GDP PAM dans les 2 groupes?

> Jpn J Thorac Cardiovasc Surg, 46 (1), 18-24 Jan 1998

## [The Effect of Pump Flow on Cerebral Oxygen Metabolism During Cardiopulmonary Bypass]

[Article in Japanese]

H Sakahashi<sup>1</sup>

2.2-2.5 L/min/m<sup>2</sup> in group L and 2.7-3.0 L/min/m<sup>2</sup> in group H

It is suggested that cerebral metabolism should be decreased during cooling to 31 degrees C of pharyngeal temperature. 2.2-2.5 l/min/m<sup>2</sup> of pump flow was adequate to keep SjO2 stable. On the other hand, it is necessary to increase pump flow to 2.7-3.0 l/min/m<sup>2</sup> during rewarming period as

**Le débit de CEC doit s'adapter à la demande métabolique**  
**variabilité du débit**





Fig. 2 Cardiac index (CI), relative mean middle cerebral arterial velocity (%MCAV) and mean arterial pressure (MAP): group L (●) and group H (■).  
%MCAV = MCAV ÷ MCAV<sub>pre-CPB</sub> × 100

## Essayer de rester physiologique = s'adapter aux besoins métaboliques



The Journal of Thoracic and Cardiovascular Surgery

Volume 134, Issue 3, September 2007, Pages 587-593



Cardiopulmonary support and physiology


## Elevated flow rate during cardiopulmonary bypass is associated with fluid accumulation

Oddbjørn Haugen MD<sup>a</sup>, Marit Farstad MD, PhD<sup>c</sup>, Venny Kvalheim MD<sup>b</sup>, Olav Bøe DDS, MSc<sup>d</sup>, Paul Husby MD, PhD<sup>e, f, g</sup>

## Type de CEC supérieur à un autre pour préserver la PAM? Limiter les contraintes vasculaires par la miniCEC?

A combination of MIECC features, such as coating, centrifugal pump, separation of cardiomy suction blood, and use of closed systems, should be considered to improve conventional CPB

Journal of Cardiothoracic and Vascular Anesthesia 35 (2021) 1–17



with ECC systems, MIECC (minimal invasive extracorporeal systems) and CCPB (conventional cardiopulmonary bypass).  
R4.4 – Il est recommandé de privilégier une technique de « CEC optimisée » afin de réduire la survenue de complications postopératoires et la mortalité hospitalière.

GRADE 1+ (accord FORT)

## Minimal invasive extracorporeal circulation (MIECC): the state-of-the-art in perfusion

Kyriakos Antoniadis, Helmut Angerer, Apostolos Dikopoulos, Polyxeni Antoniadou, Constantin Depreux, SERA University Hospital, Thessaloniki, Greece  
Christophorus H. Hoffmann, Helmut Angerer, Constantin Depreux, SERA University Hospital, 5. Stockwerk, 1. Hof in Thessaloniki, Greece  
Email: christophorus.hoffmann@sera.gr

Submitted Jan. 2019. Accepted for publication Jan. 31, 2019.






Figure 1 Established classification of MIECC circuits (3). X, pump; O, oxygenator; C, cardioplegia; T, bubble-trap/air removing device; V, vent (aortic/pulmonary); S, soft-bag/reservoir; H, hard-shell/reservoir; MIECC, minimal invasive extracorporeal circulation.

## miniCEC et PAM



If the mean arterial pressure decreased, the arterial flow was increased up to a maximum of 120% of the calculated flow. If mean arterial pressure further decreased, a 5 µg norepinephrine bolus was given to raise the MAP on CCPB between 45 and 75 mmHg.


## miniCEC et PAM: causes d'une PAM plus élevée?



### Article Efficacy of Pulsatile Flow Perfusion in Adult Cardiac Surgery: Hemodynamic Energy and Vascular Reactivity

Mikhail Dodonov <sup>1</sup>, Francesco Onorati <sup>1</sup>, Giovanni Battista Luciani <sup>1</sup>, Alessandra Franca <sup>1,\*</sup>, Maddalena Tessari <sup>1</sup>, Tiziano Menon <sup>1</sup>, Leonardo Gottin <sup>2</sup>, Aldo Domenico Milano <sup>3</sup> and Giuseppe Faggian <sup>1</sup>

- A new MEDOS Delta Stream DP3 centrifugal blood pump was used during CPB and was set in pulsatile (PP group) or non-pulsatile (NP group) mode.
- In pulsatile mode, the pump was set at maximum rotation speed variation of 3500–8500 rpm with a pulse frequency of 60 bpm; start and stop points of the pulse were set at 20 and 80% of the pulse cycle.
- This model includes the hemodynamic energy gradient as a key point rather than the pressure gradient and it is actually described in terms of energy equivalent pressure (EEP) and surplus hemodynamic energy (SHE). EEP was calculated according to the following formula:  $EEP = Rtt12 \cdot f \cdot \rho / Rtt12 \cdot f \cdot dt$  (mm Hg), where  $f$  is blood flow,  $\rho$  is blood pressure and the product of flow and pressure represents hemodynamic power



Post-op CCr, 18 h (mL/min/1.73 m <sup>2</sup> )		61 ± 35		47 ± 16 *		0.06	
		Start CPB		1h of Cross-Clamp		3h of Cross-Clamp	
		1h of Cross-Clamp		15h of Cross-Clamp		30h of Cross-Clamp	
		Re-Warming		Cross-Clamp Removal		Mean for all CPB Period	
		ANOVA P (Group/Time Interaction)					
MAP (mmHg)	PP	57 ± 15	58 ± 14	51 ± 11	58 ± 12	58 ± 14	51 ± 11
	NP	59 ± 12	61 ± 15	64 ± 19 **	69 ± 19 *	61 ± 18	57 ± 12
							56 ± 9.0
							62 ± 12 *
							0.03

## Intérêt d'une CEC pulsatile?



Contents lists available at ScienceDirect

Journal of Cardiothoracic and Vascular Anesthesia

journal homepage: [www.jcvaonline.com](http://www.jcvaonline.com)



Editorial

### Guidelines for Conduct of Cardiopulmonary Bypass

- Pulsatile perfusion may reduce postoperative pulmonary and renal complications and should be considered in patients at high risk for adverse lung and renal outcomes (grade B)

## Autres voies pour optimiser la PAM en chirurgie cardiaque?

### Closed-Loop System of Vasopressor Infusion Post Cardiac Surgery

Currently, vasopressor infusions for managing hypotension post-surgery are adjusted manually

Randomized trial of 42 patients after cardiac surgery

CLV system: Norepinephrine infusion (Two-hour study period) vs Manual (control)

Mean arterial pressure measured

% of time with hypotension (MAP < 65 mmHg): CLV - 5.4% | Control - 12.5%

% of time with hypotension (MAP 65-75 mmHg): CLV - 95% | Control - 68%

% of time with hypotension (MAP > 75 mmHg): CLV - 3.2% | Control - 28.4%

A novel closed-loop vasopressor (CLV) has been developed by physicians

Can the CLV replace manual titration?

Closed-loop vasopressor is an efficient system to control post-operative hypotension in patients with cardiac surgeries

Control of Postoperative Hypotension using a Closed-loop System for Norepinephrine Infusion in Patients after Cardiac Surgery: A Randomized Trial (Dottorini et al. 2020)



## Angiotensine II

### • PORTHOS Trial – ACTRN12623000848606

- Design: essai randomisé, en double aveugle, multicentrique, comparant une perfusion d'angiotensine II versus noradrénaline, en per-op et jusqu'à 48 h après le début de la chirurgie cardiaque sous CEC. [pmc.ncbi.nlm.nih](#)
- Population: 400 patients de chirurgie cardiaque à haut risque d'AKI (critères: Hb <130 g/L, créatinine >100 µmol/L, âge >70 ans, NYHA IV, IMC >30, au moins un critère). [pmc.ncbi.nlm.nih](#)
- Objectif: maintenir une PAM 70-80 mmHg; critère principal = durée de séjour hospitalier, critères secondaires = événements rénaux, cardiovasculaires et neurologiques. [pmc.ncbi.nlm.nih](#)
- Statut: protocole publié, essai en cours (pas encore de résultats cliniques définitifs). [pmc.ncbi.nlm.nih](#)

## Vasopressine

Randomized Controlled Trial > Am Heart J. 2024 Jun;272:86-95. doi: 10.1016/j.ahj.2024.03.008. Epub 2024 Mar 16.

### Prospective randomized double-blind study to evaluate the superiority of Vasopressin versus Norepinephrine in the management of the patient at renal risk undergoing cardiac surgery with cardiopulmonary bypass (NOVACC trial)

Pierre-Grégoire Guinot <sup>1</sup>, Olivier Desebbe <sup>2</sup>, Guillaume Besch <sup>3</sup>, Philippe Guerci <sup>4</sup>, Philippe Gaudard <sup>5</sup>, Diane Lena <sup>6</sup>, Paul Michel Mertes <sup>7</sup>, Osama Abou-Arab <sup>8</sup>, Belaid Bouhemad <sup>9</sup>; NOVACC study group

Vp mieux pour le rein?: PMID: **41208542**

## conclusion



RECOMMANDATIONS FORMALISÉES D'EXPERTS



RECOMMANDATIONS FORMALISÉES D'EXPERTS

Special Communication

**Guidelines for Perioperative Care in Cardiac Surgery**  
Enhanced Recovery After Surgery Society  
Recommendations

JAMA Surg  
Published Online May 4, 2019  
1073(5):450-755-766. doi:10.1001/jamasurg.2019.1153

Daniel T. Engelman, MD<sup>1</sup>, Waleed Ben Ali, MD<sup>2</sup>, Jason B. Williams, MD, MChD<sup>3</sup>, et al.  
> Author Affiliations | Article Information

**Réhabilitation Améliorée Après Chirurgie Cardiaque**  
**adulte sous CEC ou à cœur battant**  
ENHANCED RECOVERY AFTER CARDIAC SURGERY UNDER CPB OR OFF-PUMP

**2021**

**CHAMP 4. STRATEGIE CHIRURGICALE ET GESTION DE LA CEC**

		Avis experts	FORT
4.1	La chirurgie mitrale vidéo-assistée peut être envisagée dans des équipes entraînées		FORT
4.2	Réaliser la CEC en normothermie	1+	FORT
4.3.1	Ne pas réaliser systématiquement les pontages coronariens à cœur battant	1-	FORT
4.3.2	Les pontages coronariens à cœur battant peuvent être discutés en cas d'aorte très calcifiée	Avis experts	FORT
4.4	Utiliser une CEC optimisée	1+	FORT
4.5	Ne pas privilégier une technique de cardioplogie plutôt qu'une autre	2-	FORT

Pas de recommandations sur la PAM perCEC ou la DaO2

• Interdiscip Cardiovasc Thorac Surg. 2025 Feb 14;40(2):ivaaf002. doi: 10.1093/icvts/ivaaf002

### 2024 EACTS/EACTAIC/EBCP Guidelines on cardiopulmonary bypass in adult cardiac surgery

In summary, a high blood pressure target may result in little to no difference in patient outcomes, including AKI and deaths and further studies are needed to assess the efficacy of a higher blood pressure target among those who undergo cardiac surgery with CPB. It is more important to target DO<sub>2</sub>, optimize blood flow from CPB and monitor markers of tissue perfusion (NIRS and SvcO<sub>2</sub>) rather than use vasopressors to increase ABP (please see also section 9.7 on goal-directed perfusion).

## Conclusion

- PAM: un outil de surveillance parmi d'autre: PaCO<sub>2</sub>, débit CEC, SvO<sub>2</sub>, Hte
- La PAM est la résultante du Débit de CEC et des RVS :
  - PAM minimale si débit de CEC suffisant +++
  - Coupler la PAM à la DaO<sub>2</sub> +++ et à la demande métabolique (test de PAM sur la NIRS, sur la SVO<sub>2</sub>, pas sur l'exCO<sub>2</sub>...)
- Modifier les RVS dans un deuxième temps seulement
- Quel vasopresseur idéal?
- Titrer la PAM
  - sur organe « cible », sur microcirculation
  - maintenir PAM > 50 voire 65 mmHg sinon
  - Difficile de proposer une valeur cible de PAM selon PAM préopératoire

# Proposition personnelle d'algorithme, en normothermie

Hématocrite > 18-24%

FIO2 pour SpO2 < 99%  
(ne pas surestimer la SvO2)

Recommendations	Class <sup>a</sup>	Level <sup>b</sup>	Ref
<b>PRBC transfusions</b>			
It is recommended that PRBCs be transfused during CPB if the Hb value is <50 g/dl.	I	C	
For HCT values between 18% and 24%, PRBCs may be transfused based on an assessment of the adequacy of tissue oxygenation <sup>c</sup> .	IIb	B	[143]
PRBCs should not be transfused during CPB if the HCT is >24%.	III	C	

Débit de CEC pour SvO2 > 75%

Perfusion adaptée et non luxuriante si réflexion sur niveau de DaO2 seul

Débit de CEC pour SvO2 > 75%



PAM 40-80 mmHg



- 1 PaCO2 40-45 mmHg
- 2 PVC < 10
- 3 SpO2 < 98%
- 4 Sédation adapté (40-60 pas de BSR)

NIRS > 80% valeur de base?

oui

non

DaO2 > 300 ml/min/m<sup>2</sup>

Test noradrénaline