



Quel objectif de pression artérielle au cours de la CEC ?

DESEBBE Olivier

Département d'Anesthésie , Réanimation, et Médecine périopératoire



Clinique de la Sauvegarde, Lyon

Conflits d'intérêt

- MEDTRONIC
- BRAINDEX

Définition d'une pression

$$\text{Pascal} \longleftarrow P = \frac{F}{S} \longrightarrow \begin{array}{l} \text{Newton} \\ \text{m}^2 \end{array}$$





- Pression = Force exercée sur une surface
 - agitation incessante des molécules et de leurs collisions, entre elles ou sur des obstacles.
- transfert de quantité de mouvement dans un liquide et ses effets sur des parois.
- La pression s'exerce naturellement dans toutes les directions.

Différentes approches de la PAM

Approche électrique: loi d'OHM

- Gradient de pression motrice $\Delta P = DC \times RVS$




$$RVS = 8 \eta L / r^4$$

η = viscosité sanguine
vaisseaux

L = longueur du réseau

r = rayon des



La pression est générée par le débit et les résistances
Ex: de la CEC et l'augmentation du débit
Mettre vidéo


Histamine blockade and cardiovascular changes following heparin administration during cardiac surgery

P A Casthely ¹, D Yoganathan, B Karyanis, M Salem, T Yoganathan, C Komer, M Uribe, S Sclafani, A Hudak

Affiliations + expand

PMID: 2131900 DOI: 10.1016/s0888-6296(09)90009-6

Vasodilatation aigue




Histamine blockade and cardiovascular changes following heparin administration during cardiac surgery

P A Casthely ¹, D Yoganathan, B Karyanis, M Salem, T Yoganathan, C Komer, M Uribe, S Sclafani, A Hudak



Affiliations + expand

PMID: 2131900 DOI: 10.1016/s0888-6296(09)90009-6



The Conundrum of Systemic Arterial Pressure Management on Cardiopulmonary Bypass

[Marco Ranucci](#),* [Mauro Cotza](#), and [Umberto Di Dedda](#)




$$\Delta P = \frac{Q \times 8\eta L}{\pi R^4}$$

Heterogeneous impact of hypotension on organ perfusion and outcomes: a narrative review


Lingzhong Meng*

Department of Anesthesiology, Yale University School of Medicine, New Haven, CT, USA



E-mail: lingzhong.meng@yale.edu



Approche physiologique



- **Entrant:** Débit entrant = débit de CEC
- **Contenant:** Paroi des vaisseaux: élasticité, compliance artérielle
 - Élasticité
 - Impedance et inductance aortique (représente la resistance au flux pulsatile)
- **Contenu:** viscosité sanguine et plasmatique
- **Sortant:** Tonus vasomoteur distal: Résistances artérielles périphériques



Comment mesurer la pression artérielle?



Cathétérisme artériel et mesure invasive de la pression artérielle en anesthésie-réanimation chez l'adulte

Conférence d'experts – SFAR 1994

- Tête de pression en regard de la ligne axillaire moyenne....


Quel site permet de mesurer la PAM?

- Radial/fémoral?
- PAM radiale < PAM fémorale




Quel site permet de mesurer la PAM?

- Gradient maximal en fin de CEC et persistant
- Jusqu'à 45% des patients
 - Petite taille
 - HTA
 - procédure longue et complexe



De la PAM à la perfusion d'organe

PAM et pression de perfusion tissulaire



- 74 ans, PAC 4
- Désaturation cérébrale avec débit de CEC et PAM constantes
- Cause?
- PVC 23 mmHg....

Raisonner en terme de pression de perfusion = PAM-PVC

Pression de perfusion tissulaire

► Br J Anaesth. 2024 Jun 4;133(2):264–276. doi: [10.1016/j.bja.2024.04.046](https://doi.org/10.1016/j.bja.2024.04.046) 

PeriOperative Quality Initiative (POQI) international consensus statement on perioperative arterial pressure management

[Bernd Saugel](#)^{1,2,*}, [Nick Fletcher](#)³, [Tong J Gan](#)⁴, [Michael PW Grocott](#)⁵, [Paul S Myles](#)⁶, [Daniel I Sessler](#)⁷;

PeriOperative Quality Initiative XI (POQI XI) Workgroup Members[†], , on behalf of the

Consensus recommendation 2: We recommend increasing mean arterial pressure targets when venous or compartment pressures are elevated (strong recommendation, moderate-quality evidence).

High central venous pressure is associated with acute kidney injury and mortality in patients underwent cardiopulmonary bypass surgery

Yanli Yang¹, Jun Ma², Liyun Zhao²

Affiliations + expand

PMID: 30243200 DOI: 10.1016/j.jcrc.2018.08.034

Figure 1 Flowchart of included and excluded patients in this study.




Figure2 Curvilinear Relationship Between CVP at the end of Surgery and Morbidity of AKI




Figure3 Kaplan-Meier Analysis of Survival in two different CVP group




Patients in the high CVP group (CVP ≥ 10 cmH₂O) had a low survival rate compare to patients in low CVP group (CVP < 10 cmH₂O) by Log-Rank test (Chi-square=44.308, P<0.0001). CVP=central venous pressure.

Pour aller plus loin..... Vers la microcirculation



- Théorique
- Domaine physiologique et de recherche
- D'autres pressions sont estimées

Évolution de la Pression artérielle dans l'arbre vasculaire

- Gradient de pression selon Résistances et surface endothéliale




La PAM diminue mais la pression pulsée augmente en circulation pulsatile



Pression de perfusion capillaire

De la macrocirculation vers la microcirculation




Chute de pression dans les capillaires
Variable selon le tonus vasomoteur, les inotropes, les vasoconstricteurs

Pression intéressante = Pression d'entrée...


;

Critical Closing Pressure



Pression systémique moyenne (Pmsf)



Pression Systémique Moyenne
pression mesurée en l'absence
de flux
= Pression dans les veinules
postcapillaires (60% de la
volémie)
« obstacle aux échanges
capillaires »



zone de fermeture critique qui découple la circulation artérielle de la circulation veineuse et rend inadaptée l'utilisation de la simple PAM–PVC comme gradient de perfusion

Determination of Vascular Waterfall Phenomenon by Bedside Measurement of Mean Systemic Filling Pressure and Critical Closing Pressure in the Intensive Care Unit

Jacinta J. Maas, M.D., Rob B. de Wilde, Ph.D., Leon P. Aarts, M.D., Ph.D., Michael R. Pinsky, M.D., Dr hc, FCCM, and Jos R. Jansen, Ph.D.



The pressure gap of 26.8 ± 10.7 mmHg between Pcc and Pmsf indicates that a waterfall phenomenon is likely to be present

Monitoring of the Sublingual Microcirculation During Cardiac Surgery: Current Knowledge and Future Directions

Moritz Flick, MD^{*}, Jacques Duranteau, MD[†],
Thomas W.L. Scheeren, MD[‡], Bernd Saugel, MD^{*,§,1}

PAM et microcirculation

- ? Baisse du lactate
- Amélioration marqueurs de microcirculation?: SDF, ptiO2...
- Pas de marqueurs continus de la microcirculation disponibles pour la CEC

The response of the microcirculation to cardiac surgery



Atila Kara^{a,b}, Sakir Akin^{a,c}, and Can Ince^a

high MAP group, a higher MAP was maintained through administration of higher doses of norepinephrine and repeated bolus injections of phenylephrine. **However, there were no differences in any of the microcirculatory variables between the high and low MAP groups.**

Pression systémique per CEC

- Résultante du tonus vasomoteur et du débit de CEC
- Pression modifiable par vasodilatateurs, vasoconstricteurs, modification du nombre de tours/min

Influence du type de pompe sur la PAM en CEC



- principe de l'occlusion plus ou moins complète d'un tube souple par des galets rotatifs
- indépendantes de la postcharge: elles maintiennent leur débit quelle que soit la pression artérielle


Effet des vasoconstricteurs/vasodilatateurs sur la PAM et pas sur le débit

- turbine rotative à haute vitesse
- sensible à la précharge et à la postcharge car non occlusive

Effet mixte des vasoconstricteurs/vasodilatateurs sur la PAM et sur le débit

Test du tonus vasculaire sur la circulation systémique (DaO₂, NIRS, SvO₂, VO₂...) sera plus difficile à interpréter pour la pompe centrifuge

Quel objectif de PAM pendant la CEC?



Hypo et hypertension artérielle per CEC


- Vasoplegic syndrome during CPB may derive from
 - the release of proinflammatory cytokines: activation coagulation, inflammation
 - anaesthetic drugs
 - active endocarditis
 - preoperative use of angiotensin-converting enzyme inhibitors and calcium channel blockers.
- Hypertension during CPB may derive from
 - an inadequate level of anaesthesia/analgesia,
 - a release of catecholamines,
 - vasoconstriction due to hypothermia and others

2024 EACTS/EACTAIC/EBCP Guidelines on cardiopulmonary bypass in adult cardiac surgery

Alexander Wahba ^{1 2}, Gudrun Kunst ^{3 4}, Filip De Somer ⁵, Henrik Agerup Kildahl ^{1 2},

Avantages théoriques d'une PAM haute: 80-100


- Meilleure perfusion tissulaire (cerveau, rein, tube digestif) ;
- Meilleure adéquation aux besoins des malades hypertendus, âgés ou diabétiques ;
- Assurance d'un flux satisfaisant lorsque l'autorégulation d'un organe est perturbée ;
- Amélioration du flux collatéral pour les zones tissulaires à risque d'ischémie : « watershed stroke »




Watershed (Borderzone) Infarcts



Principles of watershed infarcts - when there is plenty all get supplied



When there is less then those at the borderzone get less



Avantages d'une PAM basse

- Moins de traumatisme pour les éléments figurés ;
- Moins d'hémorragie et de retour sanguin par les artères bronchiques
- Moins de volume aspiré et lésé dans les aspirations de cardiectomie ;
- Moins de charge embolique cérébrale ;
- Meilleure protection myocardique (réduction du flux coronarien collatéral).

Pas de vasodilatateur préférentiel

“At present, the existing literature does not contain adequate information to provide recommendations on the choice of vasodilating agents.”

Association hypotension et IRA postop

Schreiber et al. *Critical Care* (2024) 28:74
<https://doi.org/10.1186/s13054-024-04862-x>


Critical Care

Suivi 9 ans,
2300 patients

CORRESPONDENCE

Open Access

Mean arterial pressure during cardiopulmonary bypass: A modifiable risk factor for acute kidney injury in cardiac surgery patients?



b. Predicted probabilities for development of CSA-AKI with increasing TWA-MAP < 65mmHg during CPB




Fig. 1 a An example of highly granular MAP data (with one second resolution) of a patient during CPB. The AUC between the threshold

Association hypotension et complications postopératoires

5000 patients

ANESTHESIOLOGY

Intraoperative Hypotension and Acute Kidney Injury, Stroke, and Mortality during and outside Cardiopulmonary Bypass: A Retrospective Observational Cohort Study

Miguel Armengol de la Hoz, M.S.,
Valluvan Rangasamy, M.D., D.N.B., D.E.S.A.,
Andres Brenes Bastos, M.D., Xinling Xu, Ph.D.,
Victor Novack, M.D., Bernd Saugel, M.D.,
Balachundhar Subramaniam, M.D., M.P.H., F.A.S.A.

ANESTHESIOLOGY 2022; 136:927–39


EDITOR'S PERSPECTIVE

What We Already Know about This Topic

- Single-center data demonstrate that intraoperative hypotension during cardiac surgery is independently associated with stroke and acute kidney injury
- The reproducibility of this observation and whether the timing of hypotension during cardiac surgery (within vs. outside the cardiopulmonary bypass period) modifies the association remain unclear

What This Article Tells Us That Is New

- Among 4,984 patients undergoing cardiac surgery at a single tertiary care center between 2008 and 2016, 256 (5.1%) experienced the primary outcome of stroke (66, 1.3%), acute kidney injury (125, 2.5%), or mortality (109, 2.2%)
- Each 10 min of hypotension (mean arterial pressure of less than 65 mmHg) during, before, or after cardiopulmonary bypass was associated with an increased odds ratio of 1.06 (95% CI, 1.03 to 1.10; $P = 0.001$)
- Intraoperative hypotension, even if it occurs outside of cardiopulmonary bypass, is independently associated with stroke, acute kidney injury, or death after cardiac surgery



Les études: PAM haute ou basse?

Circulation

ORIGINAL RESEARCH ARTICLE



High-Target Versus Low-Target Blood Pressure Management During Cardiopulmonary Bypass to Prevent Cerebral Injury in Cardiac Surgery Patients

A Randomized Controlled Trial

**IMPROVEMENT OF
OUTCOMES AFTER
CORONARY ARTERY
BYPASS**

**A randomized trial
comparing intraoperative
high versus low mean
pressure**

Increased systemic perfusion pressure during cardiopulmonary bypass is associated with less early postoperative cognitive dysfunction and delirium[☆]

Matthias Siepe^{1,*}, Thomas Pfeiffer¹, Andreas Gieringer, Silke Zemann, Christoph Benk, Christian Schlensak, Friedhelm Beyersdorf

Department of Cardiovascular Surgery, University Medical Center Freiburg, Freiburg, Germany

Kandler et al. *Journal of Cardiothoracic Surgery* (2019) 14:107
<https://doi.org/10.1186/s13019-019-0929-4>

Journal of
Cardiothoracic Surgery

RESEARCH ARTICLE

Open Access

Higher arterial pressure during cardiopulmonary bypass may not reduce the risk of acute kidney injury



Kristian Kandler^{1*}, Jens C. Nilsson², Peter Oturai³, Mathias E. Jensen¹, Christian H. Møller¹, Jens Otto Clemmesen⁴, Henrik C. Arendrup³ and Daniel A. Steinbrüchel¹

Perfusion. 2014 Nov;29(6):496-504. doi: 10.1177/0267659114527331. Epub 2014 Mar 11.

Increasing mean arterial pressure during cardiac surgery does not reduce the rate of postoperative acute kidney injury.


Azau A¹, Markowicz P¹, Corbeau JJ¹, Cottineau C¹, Moreau X¹, Baufreton C², Beydon L³.

[+](#) Author information

Morbidité et mortalité des patients pris en charge avec des cibles élevées ou basses de pression artérielle pendant la chirurgie cardiaque sous circulation extracorporelle : une revue systématique et une méta-analyse d'études randomisées contrôlées

Charlotte C. McEwen, MD · Takhliq Amir, BHSc(Hons) · Yuan Qiu, BHSc(Hons) ·

2022



| | | n | population | Basse PAM | Haute PAM | objectif | résultats |
|---------|------|-----|---------------------|-----------|-----------|--|--|
| Azau | 2014 | 292 | À risque d'AKI | 50-60 | 75-85 | Creat > 30% | NS  |
| Siepe | 2011 | 92 | Chir coronaire | 60-70 | 80-90 | MMSE et délirium à H48 | Delirium 13% vs 0%  |
| Gold | 1995 | 248 | Chir coronaire | 50-60 | 80-100 | Complications neuro et cardiaques | 12,9 vs 4,8% 4,0 vs 1,6% décès à 6 mois  |
| Vedel | 2018 | 197 | Coronaires + valves | 40-50 | 70-80 | Volume total des lésions ischémiques cérébrales IRM POCD à J90 | NS  |
| Miao | 2022 | 40 | Valvulaire | 50-60 | 70-80 | Lactate | 3,1 versus 2,1  |
| Kandler | 2019 | 90 | PAC Valves | 40-50 | > 65 | AKI | NS  |

Increased systemic perfusion pressure during cardiopulmonary bypass is associated with less early postoperative cognitive dysfunction and delirium[☆]

Matthias Siepe^{1,*}, Thomas Pfeiffer¹, Andreas Gieringer, Silke Zemann, Christoph Benk, Christian Schlensak, Friedhelm Beyersdorf

Department of Cardiovascular Surgery, University Medical Center Freiburg, Freiburg, Germany





Fig. 2. Postoperative lactate concentration over the time on the intensive care unit. Note that the LP group's lactate rises significantly immediately after surgery (* $p < 0.05$ and ** $p < 0.001$ between the groups, error bars indicate standard error of the mean).

Lactate plus élevé dans le groupe basse PAM

Manque de données: Pas de feVG décrite, pas de PaCO₂ perCEC, pas de DaO₂/débit de CEC delirium 0% dans le groupe traité????

Table IV. Cardiac and neurologic outcomes in the two treatment groups (intention to treat)

| | <i>Low MAP</i> (<i>n</i> = 124) | | <i>High MAP</i> (<i>n</i> = 124) | | <i>Low – high MAP</i> | | |
|---|-------------------------------------|----------|--------------------------------------|----------|-----------------------|----------|------------------------------------|
| | <i>No.</i> | <i>%</i> | <i>No.</i> | <i>%</i> | <i>No.</i> | <i>%</i> | <i>95% CI for % difference</i> |
| Fatal stroke | 2 | 1.6 | 0 | 0.0 | | | |
| Hemiparesis* | 2 | 1.6 | 1 | 0.8 | | | |
| Aphasia | 3 | 2.4 | 1 | 0.8 | | | |
| Cortical blindness | 1 | 0.8 | 0 | 0.0 | | | |
| Monocular blindness | 1 | 0.8 | 0 | 0.0 | | | |
| Other focal deficit | 0 | 0.0 | 1 | 0.8 | | | |
| Total permanent neurologic complications | 9 | 7.2 | 3 | 2.4 | 6 | 4.8 | –0.5, 11.0 |
| Fatal cardiogenic shock | 1 | 0.8 | 2 | 1.6 | | | |
| Shock | 1 | 0.8 | 0 | 0.0 | | | |
| Myocardial infarction | 4 | 3.2 | 1 | 0.8 | | | |
| Total cardiac complications | 6 | 4.8 | 3 | 2.4 | 3 | 2.4 | –2.2, 7.1 |
| Other death, total (not attributable to cardiac or neurologic causes) | 2 | 1.6 | 0 | 0 | 2 | 1.6 | –0.6, 3.8 |
| Total mortality and major cardiac and neurologic morbidity | 16† | 12.9 | 6 | 4.8 | 10 | 8.1 | 1.0, 15.1 |

Taux d'AIC dans le groupe low MAP très élevé et grave
Débit de CEC 1,9 à 2,3 l/min/m²

RESEARCH


Open Access



Target blood pressure management during cardiopulmonary bypass improves lactate levels after cardiac surgery: a randomized controlled trial

Qing Miao¹, Dong Jin Wu¹, Xu Chen¹, Meiyong Xu¹, Lin Sun², Zhen Guo², Bin He³ and Jingxiang Wu^{1*}

- Débit de CEC 2,4 l/min/m², hypothermie modérée




RESEARCH

Open Access



Target blood pressure management during cardiopulmonary bypass improves lactate levels after cardiac surgery: a randomized controlled trial

Qing Miao¹, Dong Jin Wu¹, Xu Chen¹, Meiyong Xu¹, Lin Sun², Zhen Guo², Bin He³ and Jingxiang Wu^{1*}



Cr (μM·L⁻¹)

Baseline

78 [63, 90]

74 [60, 84]

0.558

6 h after the operation

72 [58.8, 88]

79 [52, 93]

0.690

24 h after the operation

75 [57, 91]

75 [63, 86]

0.706

RI IN (mM·L⁻¹)



High-Target Versus Low-Target Blood Pressure Management During Cardiopulmonary Bypass to Prevent Cerebral Injury in Cardiac Surgery Patients
A Randomized Controlled Trial

Table 1. Baseline Characteristics

| | Low-Target Group (n=99) | High-Target Group (n=98) |
|---|-------------------------|--------------------------|
| Age, y | 65.0±10.7 | 69.4±8.9 |
| Male sex, n (%) | 93 (93.9) | 84 (85.7) |
| Nonwhite race, n (%) | 2 (2.0) | 0 (0) |
| Previous myocardial infarction, n (%)* | 37 (37.4) | 37 (37.8) |
| Recent myocardial infarction (past 2 wk), n (%) | 30 (30.3) | 25 (25.5) |
| Aortic valvular disease, n (%) | 34 (34.3) | 34 (34.7) |
| Angina, CCS score >1, n (%)† | 64 (64.6) | 47 (48.0) |
| Current or previous atrial fibrillation, n (%) | 14 (14.1) | 13 (13.3) |
| Hypertension, n (%) | 83 (84.8) | 87 (88.8) |
| Diabetes mellitus, type 1 or 2 (insulin treated), n (%) | 10 (10.1) | 10 (10.2) |
| Diabetes mellitus, type 2 (non–insulin treated), n (%) | 14 (14.1) | 14 (14.3) |
| Chronic lung disease, n (%) | 9 (9.1) | 12 (12.2) |
| Current smoker, n (%) | 18 (18.2) | 15 (15.3) |
| Current alcohol abuse, n (%) | 7 (7.1) | 7 (7.1) |
| BMI, kg/m ² ‡ | 27.0 (3.8) | 27.6 (4.0) |
| Left ventricular ejection fraction, n (%) | | |
| >50% | 54 (54.5) | 50 (51.5) |



High-Target Versus Low-Target Blood Pressure Management During Cardiopulmonary Bypass to Prevent Cerebral Injury in Cardiac Surgery Patients

A Randomized Controlled Trial

| | Low-Target Group (n=98) | High-Target Group (n=97) |
|--|--------------------------------|---------------------------------|
| MAP before anesthesia induction, mm Hg | 92.3±15.7 | 96.9±13.4 |
| MAP during bypass, mm Hg | 44.7±4.7 | 66.8±4.9 |
| Norepinephrine infused in the OR, µg/kg | 2.65±6.01 | 17.43±20.14 |
| Patients receiving norepinephrine in the OR, n (%) | 35 (35.7) | 90 (92.7) |

High-Target Versus Low-Target Blood Pressure Management During Cardiopulmonary Bypass to Prevent Cerebral Injury in Cardiac Surgery Patients

A Randomized Controlled Trial

Overall, diffusion-weighted imaging revealed new cerebral lesions in 52.8% of patients in the low-target group versus 55.7% in the high-target group ($P=0.76$). The primary outcome of volume of new cerebral lesions was

| | Low-Target Group, n | | High-Target Group, n | | Difference (95% CI) | OR (95% CI) | P Value |
|---|---------------------|------------|----------------------|------------|------------------------------|-------------|-------------------|
| Complete cases, mean (SD) Volume AVC | 89 | 415 (2682) | 80 | 488 (2539) | 8 (-978 to 994) [†] | | 0.99 [‡] |
| Adverse events, n (%) | | | | | | | |
| Death | 99 | 0 (0) | 98 | 4 (4.1) | | Inf | 0.06 |
| Creatinine, doubling of baseline value, n (%) | 2 (2.0) | | 9 (9.4) | | 4.93 (1.02–48.12) | 0.03 | |

- PAM basse et moins d'AVC , moins d'insuffisance rénale

Conclusions

No difference in mortality nor in the level of cognitive functioning was found according to blood pressure target during cardiac surgery long-term at 3-year follow-up.

RESEARCH ARTICLE

Open Access



Higher arterial pressure during cardiopulmonary bypass may not reduce the risk of acute kidney injury

Kristian Kandler^{1*}, Jens C. Nilsson², Peter Oturai³, Mathias E. Jensen¹, Christian H. Møller¹, Jens Otto Clemmesen⁴, Henrik C. Arendrup¹ and Daniel A. Steinbrüchel¹

38% groupe PAM 40-50
versus
46% groupe PAM > 65

90 patients

Table 3 Postoperative data

| | CG | HPG | p value | 95% CI |
|------------------------------------|------------|------------|---------|-----------|
| Delta sCr (mmol/L) | 20 (6–51) | 25 (10–49) | 0.560 | –18 to 18 |
| Delta eGFR (ml/min) | –6 (–15–4) | –3 (–13–5) | 0.522 | –7 to 5 |
| AKI | 16 (38%) | 19 (46%) | 0.447 | |
| Dialysis | 4 (10%) | 3 (7%) | 0.565 | |
| Re-operation | 4 (10%) | 1 (3%) | 0.165 | |
| Stroke | 3 (8%) | 2 (5%) | 0.687 | |
| 4 months follow-up | | | | |
| Change in GFR (ml/min) | –9 ± 12 | –5 ± 16 | 0.288 | –13 to 4 |
| > 10% decrease in GFR ^a | 11 (44%) | 9 (39%) | 0.732 | |
| 30-day mortality | 3 (7%) | 3 (7%) | 1.000 | |
| 6-months mortality | 7 (16%) | 8 (18%) | 0.763 | |

Percentages are given as total within group. Continuous data are presented as mean ± SD or median (interquartile min – max)

AKI Acute kidney injury, CI confidence interval, AP Arterial pressure, eGFR Estimated glomerular filtration rate (calculated using the Cockcroft-Gault formula), GFR Glomerular filtration rate (estimated by Cr-EDTA clearance), sCr Serum creatinine

^aPercentages of total of patients at follow-up

Prévention de l'AKI par la PAM?

■ SYSTEMATIC REVIEW ARTICLE

Society of Cardiovascular Anesthesiologists Clinical Practice Update for Management of Acute Kidney Injury Associated With Cardiac Surgery

Ke Peng, MD, PhD,*† David R. McLroy, MBBS,‡ Bruce A. Bollen, MD,§ Frederic T. Billings IV, MD,‡ Alexander Zarbock, MD,|| Wanda M. Popescu, MD,¶ Amanda A. Fox, MD, MPH,# Linda Shore-Lesserson, MD,** Shaofeng Zhou, MD,†† Mariya A. Geube, MD, ‡‡ Fuhai Ji, MD, PhD,† Meena Bhatia, MD,§§ Nanette M. Schwann, MD,||| Andrew D. Shaw, MB, FCCM, FFICM, FRCA,¶¶ and Hong Liu, MD, FASE*

■ SYSTEMATIC REVIEW ARTICLE

| Table 1. Study Characteristics | | | | | | |
|---|---------|---------------------------------|-----|---|--------------------------|----------------------------------|
| Study | Region | Procedure | N | Interventional and control treatments, and main findings on CS-AKI | AKI definition | Journal (IF2020) |
| Intraoperative target blood pressure | | | | | | |
| Azau et al 2014 ¹¹ | France | CABG, valve, or aorta, with CPB | 292 | A high level of MAP (75–85 mm Hg) versus control (MAP 50–60 mm Hg) during normothermic CPB did not reduce the risk of AKI. | 30% rise in sCr | <i>Perfusion</i> (1.97) |
| Kandler et al 2019 ¹² | Denmark | CABG + valve, with CPB | 90 | Arterial pressure >60 mmHg versus control (MAP 47 mm Hg) during CPB did not reduce the incidence of AKI or chronic kidney injury at postoperative 4 mo. | RIFLE | <i>J Cardiothorac Sur</i> (1.64) |
| Vedel et al 2018 ¹³ | Denmark | CABG and/or valve, with CPB | 197 | A higher MAP (70–80 mm Hg) versus control (MAP 40–50 mm Hg) increased the number of patients with postoperative doubling of sCr. | sCr ≥2 times of baseline | <i>Circulation</i> (29.7) |

Métaanalyse 2022

| Outcome (number of studies) | High MAP target, <i>n</i> /total <i>N</i> | Low MAP target, <i>n</i> /total <i>N</i> | Relative risk | 95% CI | <i>P</i> value | <i>I</i> ² (%) |
|------------------------------|---|--|--------------------------|-------------|----------------|---------------------------|
| Dichotomous variables | | | | | | |
| Hospital mortality (4) | 8/311 | 6/316 | 1.1 | 0.4 to 3.3 | 0.84 | 0 |
| 30-day mortality (4) | 12/334 | 6/337 | 1.6 | 0.6 to 4.36 | 0.33 | 0 |
| 6-month mortality (3) | 15/310 | 19/309 | 0.8 | 0.4 to 1.6 | 0.55 | 0 |
| Myocardial infarction (3) | 7/367 | 8/367 | 0.9 | 0.3 to 3.2 | 0.89 | 18 |
| Delirium (2) | 10/142 | 13/147 | 0.5 | 0.03 to 8.6 | 0.61 | 75 |
| Cognitive decline (2) | 39/190 | 35/204 | 1.2 | 0.7 to 2.1 | 0.46 | 35 |
| Stroke (3) | 11/261 | 13/266 | 1.0 | 0.2 to 4.9 | 0.95 | 65 |
| Hemodialysis (5) | 11/368 | 11/372 | 1.0 | 0.5 to 2.3 | 0.98 | 0 |
| RIFLE “risk” criteria (4) | 120/297 | 98/303 | 1.3 | 0.9 to 1.8 | 0.25 | 41 |
| Number transfused (3) | 85/227 | 60/229 | 1.4 | 1.1 to 1.9 | 0.004 | 0 |
| Outcome (number of studies) | High MAP target, mean (SD) | Low MAP target, mean (SD) | Weighted mean difference | 95% CI | <i>P</i> value | <i>I</i> ² (%) |
| Continuous variables | | | | | | |
| ICU LOS in days (4) | 2.4 (4.7) | 2.6 (5.3) | 0.2 | -0.4 to 0.8 | 0.55 | 29 |
| Hospital LOS in days (4) | 10.7 (7.8) | 11.1 (9.4) | 1.1 | 0.3 to 1.8 | 0.006 | 13 |
| PRBCs transfused/patient (3) | 2.1 (2.7) | 1.9 (2.9) | 0.1 | -0.1 to 0.3 | 0.35 | 0 |

CI = confidence interval; ICU = intensive care unit; MAP = mean arterial pressure; LOS = length of stay; PRBC = packed red blood cell; RIFLE = Risk, Injury, Failure, Loss, and End-stage Kidney classification

Review > [Cochrane Database Syst Rev. 2022 Nov 30;11\(11\):CD013494.](#)

doi: [10.1002/14651858.CD013494.pub2](https://doi.org/10.1002/14651858.CD013494.pub2).

High versus low blood pressure targets for cardiac surgery while on cardiopulmonary bypass

[Yuki Kotani](#)¹, [Yuki Kataoka](#)^{2 3 4 5}, [Junichi Izawa](#)^{6 7}, [Shoko Fujioka](#)⁸, [Takuo Yoshida](#)^{9 10},
[Junji Kumasawa](#)^{11 12}, [Joey Sw Kwong](#)¹³

Affiliations + expand

PMID: 36448514 PMCID: PMC9709767 (available on 2023-11-30)

DOI: [10.1002/14651858.CD013494.pub2](https://doi.org/10.1002/14651858.CD013494.pub2)

Conclusions des auteurs

Une cible de pression artérielle élevée pourrait n'entraîner que peu ou pas de différence dans les critères de jugement relatifs aux patients, y compris les lésions rénales aiguës et la mortalité. Compte tenu de largeur des intervalles de confiance, d'autres études sont nécessaires pour confirmer l'efficacité d'une cible de pression artérielle plus élevée chez les personnes qui subissent une chirurgie cardiaque avec pontage cardio-pulmonaire.

Reco EACTS 2024 sur la PAM per CEC

Recommendation Table 29 Recommendations for kidney protection

| Recommendations | Class ^a | Level ^b | Ref ^c |
|--|--------------------|--------------------|---------------------------------|
| Routinely targeting a high MAP using vasoconstrictors is not recommended during CPB to reduce AKI. | III | A | 377,378,380,381 |

Reco EACTS 2024



Recommendation Table 40 Recommendations for control of mean arterial blood pressure during cardiopulmonary bypass

| Recommendations | Class ^a | Level ^b | Ref ^c |
|--|--------------------|--------------------|------------------|
| It is recommended that the MAP be maintained between 50 and 80 mmHg with vasoconstrictors and vasodilators if required, having ensured that the depth of anaesthesia and pump flow rate are sufficient. | I | A | 381,511 |
| The use of vasopressors to increase the MAP to values above 80 mmHg during CPB is not recommended. | III | B | 381,510,517 |
| Targeting the MAP during CPB within the limits of individualized cerebral autoregulation data, measured under normocapnic conditions before CPB, should be considered whenever the technical and human skills are available. | IIa | A | 222,519,520 |
| It is recommended that vasoplegic syndrome during CPB be treated with α 1-adrenergic agonists and/or vasopressin. | I | C | 521,523 |
| In refractory vasoplegic syndrome, alternative drugs (methylene blue or terlipressin) should be considered, alone or in combination. | IIa | B | 522,523 |
| Hydroxocobalamin or angiotensin II may be considered to treat vasoplegic syndrome during CPB. | IIb | C | 524–527 |

Titre la PAM pendant la CEC?

= individualiser pour chaque patient le niveau de PAM selon ses besoins


Détermination physiologique de la PAM idéale



Optimisation de la PAM? Autorégulation de l'organe

Pas de variation du DSR et du DFG pour des variations de pression artérielle moyenne entre 80 et 180 mmHg

Cerveau
Rein
Myocarde (avant/après clampage)
Mésentère?



Titration selon la valeur initiale de PAM?

**Improvement of Outcomes after
Coronary Artery Bypass II: A
Randomized Trial Comparing
Intraoperative High Versus Customized
Mean Arterial Pressure**

Mary E. Charlson, M.D.,* Janey C. Peterson, Ed.D., R.N.,* Karl H. Krieger, M.D.,†

Adaptation PAM per CEC selon valeur basale PAM

- Sédation, pose KT artériel, attente 4 min
- the next 6 mean blood pressures downloaded to the computer from the operating room monitor (at 10 sec intervals) were recorded and an average MAP was taken.

Improvement of Outcomes after Coronary Artery Bypass II: A Randomized Trial Comparing Intraoperative High Versus Customized Mean Arterial Pressure

Mary E. Charlson, M.D.,* Janey C. Peterson, Ed.D., R.N.,* Karl H. Krieger, M.D.,†

- **412 patients**
- **Chir coronaire programmée**
- **80 +/- 20 mmHg,**
- **versus**
- **max 90 (57-90), 88 moy**
- **Complications neuro et
cardiaques/cognitives: NS 16,5 vs
14,5%**
- Among those whose average MAP was more than 20 mmHg below their autoregulatory range, the cardiac and neurologic complication rate was double that of patients whose MAP was not that low (15.9% vs. 7.3%).
- 61% adhérence gp PAM préop
- 85% adhérence gp Pam 80
- management at higher MAP on CPB dramatically reduces stroke occurrence in patients with severe aortic atheromatous disease (grade IV and V)

**Improvement of Outcomes after
Coronary Artery Bypass II: A
Randomized Trial Comparing
Intraoperative High Versus Customized
Mean Arterial Pressure**

Mary E. Charlson, M.D.,* Janey C. Peterson, Ed.D., R.N.,* Karl H. Krieger, M.D.,†

Même PAM dans les 2 groupes...

| B. Intraoperative Management | | | |
|-------------------------------------|-----------------|-------------------|----------|
| Pre-bypass | High MAP | Custom MAP | P |
| Pre-bypass MAP (mmHg) | 86 ± 8 | 85 ± 9 | |
| Pre-bypass cardiac output (L/min) | 4 ± 1 | 4 ± 1. | |
| Bypass Times (min) | | | |
| Bypass | 74 ± 24 | 77 ± 22 | ns |
| Cross-clamp | 40 ± 15 | 40.4 ± 14 | ns |
| Pump flows (L/min/m ²)# | | | |
| Bypass on—warming (± θ) | 2.0 ± 0.3 | 2.0 ± 0.3 | ns |
| Warming—cross-clamp off (± θ) | 2.1 ± 0.3 | 2.2 ± 0.3 | ns |
| Cross-clamp off—bypass off (± θ) | 2.4 ± 0.2 | 2.34 ± 0.3 | ns |

observationnel

Difference between pre-operative and cardiopulmonary bypass mean arterial pressure is independently associated with early cardiac surgery-associated acute kidney injury

Hussein D Kanji, Costas J Schulze, Marilou Hervas-Malo, Peter Wang, David B Ross, Mohamad Zibdawi and Sean M Bagshaw ✉

Journal of Cardiothoracic Surgery 2010 5:71

<https://doi.org/10.1186/1749-8090-5-71> | © Kanji et al; licensee BioMed Central Ltd. 2010

Received: 7 May 2010 | **Accepted:** 8 September 2010 | **Published:** 8 September 2010

Nonpulsatile pump flow rates were kept at 2.4 L/min/m₂

Delta MAP > 26 mmHg

Mesurer en site fémoral si petit gabarit, procédure longue

Position de la tête de pression...

Difference between pre-operative and cardiopulmonary bypass mean arterial pressure is independently associated with early cardiac surgery-associated acute kidney injury

Table 4 Multi-variable adjusted logistic regression model[†] of association between delta MAP and CSA-AKI

| Parameter | Odds Ratio | 95% CI | P-value |
|-------------------------------------|------------|----------|---------|
| Male sex | 0.7 | 0.3-1.7 | 0.49 |
| Age ≥75 years (present) | 2.1 | 0.9-4.9 | 0.08 |
| BMI ≥25 kg/m ² (present) | 4.2 | 1.6-11.2 | 0.0039 |
| Delta MAP ≥26 mmHg (present) | 2.8 | 1.3-6.1 | 0.009 |
| Flow ≥54 per mL/kg/min (present) | 0.3 | 0.1-0.7 | 0.004 |
| Side-biting clamp (present) | 3.0 | 1.3-7.1 | 0.012 |

Abbreviations: BMI = Body Mass Index; MAP = mean arterial pressure; CPB = cardiopulmonary bypass

Model characteristics: C-statistic = 0.788



ANESTHESIOLOGY


Automated Ambulatory Blood Pressure Measurements and Intraoperative Hypotension in Patients Having Noncardiac Surgery with General Anesthesia

A Prospective Observational Study

Bernd Saugel, M.D., Philip C. Reese, M.D.,
Daniel I. Sessler, M.D., Christian Burfeindt,
Julia Y. Nicklas, M.D., Hans O. Pinnschmidt, Ph.D.,
Daniel A. Reuter, M.D., Stefan Südfeld, M.D.

ANESTHESIOLOGY 2019; 131:74–83

Nighttime, lowest value



ANESTHESIOLOGY

Automated Ambulatory Blood Pressure Measurements and Intraoperative Hypotension in Patients Having Noncardiac Surgery with General Anesthesia

A Prospective Observational Study

Bernd Saugel, M.D., Philip C. Reese, M.D.,
Daniel I. Sessler, M.D., Christian Burfeindt,
Julia Y. Nicklas, M.D., Hans O. Pinnschmidt, Ph.D.,
Daniel A. Reuter, M.D., Stefan Südfeld, M.D.

ANESTHESIOLOGY 2019; 131:74–83





Fig. 6. Difference between lowest nighttime mean arterial pressure (MAP) and a MAP of 65 mmHg. Histogram showing the frequency (n; y axis; n total = 370) of the difference between the lowest nighttime MAP and a MAP of 65 mmHg. The median difference (25th to 75th percentile range) was +5 (–1 to +13) mmHg. The *dotted vertical line* represents the median difference.

Ajuster la PAM selon la circulation
cérébrale?

DSC



Upper Limit of Autoregulation
ULA


Lower Limit of Autoregulation
LLA

63-71

84-90

PAM

DSC



Upper Limit of Autoregulation
ULA

Lower Limit of Autoregulation
LLA

Mx = 1

Mx = 0,4

Mx = 0

Mx = 0,4

Mx = 1


Enregistrement continu et simultané
Analyse sur logiciel dédié de paires de
données PAM/DSC
Corrélation de Pearson sur une période de
temps (5 min/30 min): **Mean flow index =
Mx**

63-71

84-90

PAM


DSC



Enregistrement continu
Analyse sur logiciel dédié
données PAM/DSC
Corrélation de Pearson
temps (5 min/30 min):
Mx

PAM

A




Calcul rapide
Mx

Calcul
historique
du Mx

Valeur du Mx selon le
niveau de PAM du
patient





PAM
 Research Software - Not for medical usage.



Optimal blood pressure during cardiopulmonary bypass defined by cerebral autoregulation monitoring



Daijiro Hori, MD,^a Yohei Nomura, MD,^a Masahiro Ono, MD,^b Brijen Joshi, MD,^c Kaushik Mandal, MD,^a Duke Cameron, MD,^a Masha Kocherginsky, PhD,^d and Charles W. Hogue, MD^e




FIGURE 2. Representative graph of cerebral autoregulation monitoring by Mx during CPB. The top graph represents the time-series of ABP, and the bottom bar-graph represents the percentage of the time of the recording spent at 5 mm Hg bin. Optimal MAP (ABP) for the left and right side of the brain was defined as that MAP with the lowest Mx. LLA and ULA were defined as the MAP at which Mx reached 0.4. In this example, the optimal MAP is 85 mm Hg (*black arrow*) and the LLA and ULA are 70 mm Hg and 95 mm Hg, respectively (*black-outlined arrow*). *ABP*, Arterial blood pressure.

Variabilité individuelle de la LLA

Predicting the Limits of Cerebral Autoregulation During Cardiopulmonary Bypass

Brijen Joshi, MD,* Masahiro Ono, MD,† Charles Brown, MD,* Kenneth Brady, MD,‡ R. Blaine Easley, MD,§ Gayane Yenokyan, PhD,|| Rebecca F. Gottesman, MD, PhD,¶ and Charles W. Hogue, MD*




Figure 1. Number of subjects versus the mean arterial blood pressure at the lower limit of cerebral blood flow autoregulation during cardiopulmonary bypass based on the transcranial Doppler-determined mean velocity index.

Détermination du LLA avec l'oxymétrie cérébrale




Figure 5. Distributions of LLAs—cotrending algorithm and TCD reference. A, Bland-Altman plot of the data displaying mean bias = 0.22 mmHg, an upper LOA = 10.79 mmHg (95% CI, 10.09–11.49) and a lower LOA = -10.35 mmHg (95% CI, -9.65 to -11.05). B, Scatter plot of individual LLAs over time over all cases (bubble sizes proportional to number of collocated data points). The dashed line in the middle represents the unity line, which indicates a perfect agreement between the method and the reference. CI indicates confidence interval; LLA, lower limit of autoregulation; LOA, limit of agreement.

Les études observationnelles

Blood Pressure Excursions Below the Cerebral Autoregulation Threshold During Cardiac Surgery Are Associated With Acute Kidney Injury


Masahiro Ono, MD, PhD¹, George J. Arnaoutakis, MD¹, Derek M. Fine, MD², Kenneth Brady,

Published in final edited form as:

J Thorac Cardiovasc Surg. 2014 January ; 147(1): . doi:10.1016/j.jtcvs.2013.07.069.

Duration and magnitude of blood pressure below cerebral autoregulation threshold during cardiopulmonary bypass is associated with major morbidity and operative mortality


Masahiro Ono, MD, PhD^a, Kenneth Brady, MD^b, R. Blaine Easley, MD^b, Charles Brown,



| Variable | No MMOM (n = 354) | MMOM (n = 83) | P value |
|---|-------------------------|-------------------------|---------|
| Average MAP during CPB (mm Hg) | 74 ± 8 (73–75) | 75 ± 9 (72–76) | .203 |
| Average rScO ₂ | 54 ± 11 (52–55) | 55 ± 7 (53–56) | .388 |
| Average COx | 0.27 ± 0.18 (0.25–0.29) | 0.26 ± 0.17 (0.21–0.29) | .749 |
| LLA (mm Hg) | 69 ± 14 (67–70) | 71 ± 12 (67–72) | .136 |
| AUC _{MAP<LLA} (mm Hg × min/h) | 2.4 (1.1–5.7) | 6.5 (2.1–15.4) | .017 |

Arterial pressure above the upper cerebral autoregulation limit during cardiopulmonary bypass is associated with postoperative delirium

D. Hori¹, C. Brown², M. Ono¹, T. Rappold², F. Sieber², A. Gottschalk², K. J. Neufeld³, R. Gottesman⁴, H. Adachi⁵ and C. W. Hogue^{2*}



2 études prospectives

JAMA Surgery | **Original Investigation**

Effect of Targeting Mean Arterial Pressure During Cardiopulmonary Bypass by Monitoring Cerebral Autoregulation on Postsurgical Delirium Among Older Patients A Nested Randomized Clinical Trial

Charles H. Brown IV, MD, MHS; Karin J. Neufeld, MD, MPH; Jing Tian, MS; Julia Probert, BA; Andrew LaFlam, BA; Laura Max, MHS, PA-C; Daijiro Hori, MD; Yohei Nomura, MD; Kaushik Mandal, MD; Ken Brady, MD; Charles W. Hogue, MD; and the Cerebral Autoregulation Study Group

Randomized Controlled Trial > [Semin Thorac Cardiovasc Surg.](#) 2021 Summer;33(2):429-438.


doi: 10.1053/j.semtcvs.2020.09.032. Epub 2020 Nov 10.

Personalized Blood Pressure Management During Cardiac Surgery With Cerebral Autoregulation Monitoring: A Randomized Trial

Charles W Hogue ¹, Charles H Brown 4th ², Daijiro Hori ³, Masa Ono ⁴, Yohei Nomura ³,
Lauren C Balmert ⁵, Nina Srdanovic ⁵, Jordan Grafman ⁶, Kenneth Brady ⁷,
Cerebral Autoregulation Study Group

Affiliations + expand

PMID: 33186735 DOI: 10.1053/j.semtcvs.2020.09.032



The lower limit of autoregulation was determined by the senior author (C.W.H.) before CPB based on the highest MAP where Mx increased from less than 0.4 to 0.4 or greater.¹


When Mx did not cross 0.4 clearly, the lower limit of autoregulation was defined as the blood pressure with the lowest Mx (the MAP with the best autoregulation)

Effect of Targeting Mean Arterial Pressure During Cardiopulmonary Bypass by Monitoring Cerebral Autoregulation on Postsurgical Delirium Among Older Patients

A Nested Randomized Clinical Trial

Charles H. Brown IV, MD, MHS; Karin J. Neufeld, MD, MPH; Jing Tian, MS; Julia Probert, BA; Andrew LaFlam, BA; Laura Max, MHS, PA-C; Daijiro Hori, MD; Yohei Nomura, MD; Kaushik Mandal, MD; Ken Brady, MD; Charles W. Hogue, MD; and the Cerebral Autoregulation Study Group

Figure 2. Delirium Incidence by Randomization Group



Mean arterial pressure during cardiopulmonary bypass was managed to standard care or autoregulation-targeted goals. Delirium incidence randomization group is shown.

Table 2. Characteristics of Management During Cardiopulmonary Bypass for Patients Randomized to Standard Care vs Autoregulation-Targeted Management of Mean Arterial Pressure

| Characteristic of Management | Standard Care (n = 94) | Autoregulation-Targeted (n = 105) | P Value |
|--|------------------------|-----------------------------------|---------|
| Phenylephrine, median (IQR), mg | 1.2 (0.3-2.3) | 1.8 (0.5-3.6) | .02 |
| Vasopressin administration, No. (%) | 6 (6.4) | 9 (8.6) | .56 |
| Cardiopulmonary bypass flow, mean (SD), L/min | 4.4 (0.6) | 4.4 (0.6) | .92 |
| Isoflurane, mean (SD), % | 0.76 (0.27) | 0.77 (0.31) | .71 |
| Arterial pressure during cardiopulmonary bypass, mean (SD), mm Hg | 71.3 (7.6) | 73.9 (6.7) | .01 |
| Arterial pressure at the lower limit of autoregulation, mean (SD), mm Hg | 68.7 (11.3) | 66.0 (10.9) | .10 |
| Product of the duration of time and mean arterial pressure below the lower limit of autoregulation, median (IQR), mm Hg × h ^a | 9.5 (3.7-19.5) | 5.3 (2.0-13.4) | .002 |


ADULT – Original Submission

Personalized Blood Pressure Management During Cardiac Surgery With Cerebral Autoregulation Monitoring: A Randomized Trial

Charles W. Hogue MD ^{*}, Charles H. Brown IV MD, MPH [†], Daijiro Hori MD [‡], Masa Ono MD [§], Yohei Nomura MD [‡], Lauren C. Balmert PhD [¶], Nina Srdanovic MS [¶], Jordan Grafman PhD ^{||}, Kenneth Brady MD ^{**}, The Cerebral Autoregulation Study Group [#]

Show more

AVC, troubles cognitifs



a reduction in the frequency of delirium and better performance on tests of memory 4-6 weeks after surgery

Limites de
l'approche
de la PAM
titrée sur la
circulation
cérébrale

Hemodilution Combined With Hypercapnia Impairs Cerebral Autoregulation During Normothermic Cardiopulmonary Bypass

Ervin E. Ševerdija, EKP,* Nousjka P.A. Vranken, BAsc,* Antoine P. Simons, EKP, PhD,* Erik D. Gommer, MSc, PhD,† John H. Heijmans, MD, PhD,‡ Jos G. Maessen, MD, PhD,* and Patrick W. Weerwind, CCP, PhD*





Fig 1. Cerebral autoregulation as indicated by the autoregulation index (ARI) at baseline, at 3 levels of p_aCO₂, and at 2 levels of hematocrit during cardiopulmonary bypass.


Effet de la PaCO₂ et de la PaO₂ sur le DSC



Altération de l'AC et réchauffement....

Joshi et al.

Page 11



Published in final edited form as:

Anesth Analg. 2010 February 01; 110(2): 321–328. doi:10.1213/ANE.0b013e3181c6fd12.

Impaired Autoregulation of Cerebral Blood Flow During Rewarming from Hypothermic Cardiopulmonary Bypass and Its Potential Association with Stroke

Brijen Joshi, MD*, Kenneth Brady, MD*, Jennifer Lee, MD*, Blaine Easley, MD*, Rabi

Table 5

Neurological Outcomes for Patients with and Without Impaired Cerebral Blood Flow Autoregulation During Rewarming on Cardiopulmonary Bypass

| Outcome | No impairment (n = 60) | Impairment (n = 67) | P |
|---------------------------|------------------------|---------------------|-------|
| Perioperative stroke | 0 | 7 (10.4%) | 0.015 |
| Transient ischemic attack | 1 (1.7%) | 0 | 0.463 |

Nonpulsatile CPB with a nonocclusive roller pump was used, and CPB flow was maintained between 2.0 and 2.4 L · min⁻¹ · m⁻².

Limites de l'optimisation de la PAM sur le doppler TC

- Technique à apprendre: ACM?, position
- Matériel spécifique: connecter en continu le signal de PAM et le doppler/NIRS
- LLA évolutive au cours du temps et des conditions: température, pulsatilité, PaCO₂, Hte,

Études en cours

Trial to Compare Different Strategies of Mean Arterial Pressure Management During Cardiopulmonary Bypass

I Integrated University Hospital Trust of Verona

Status

Enrolling

Randomized Controlled Trial > Am Heart J. 2023 Jul;261:10-20. doi: 10.1016/j.ahj.2023.03.005.

Epub 2023 Mar 18.

Perioperative individualized hemodynamic optimization according to baseline mean arterial pressure in cardiac surgery patients: Rationale and design of the OPTIPAM randomized trial

Richard Descamps¹, Julien Amour², Emmanuel Besnier³, Adrien Bougle⁴, H  l  ne Charbonneau⁵, Martin Charvin⁶, Bernard Cholley⁷, Olivier Desebbe⁸, Jean-Luc Fallahi⁹, Denis Frasca¹⁰, Fran  ois Labaste¹¹, Elane Lena¹², Yazine Mahjoub¹³, Paul-Michel Meentes¹⁴, Serge Mollieux¹⁵, Pierre-Henri Moury¹⁶, Mouhamed Djahoum Moussa¹⁷, Jean-Ferreol Oilleau¹⁸, Alexandre Ouattara¹⁹, Sophie Provenchere²⁰, Bertrand Rozec²¹, Jean-Jacques Parienti²², Marc-Olivier Fischer²³; OPTIPAM investigators

> Trials. 2024 Mar 15;25:191. doi: [10.1186/s13063-024-07992-3](https://doi.org/10.1186/s13063-024-07992-3)

Mean Arterial Pressure (MAP) Trial: study protocol for a multicentre, randomized, controlled trial to compare three different strategies of mean arterial pressure management during cardiopulmonary bypass

Alessandra Francica^{1,✉}, Gina Mazzeo¹, Antonella Galeone¹, Daniele Linardi¹, Livio San Biagio¹, Giovanni Battista Luciani¹, Francescoonorati¹

> Author information > Article notes > Copyright and License information

PMCID: PMC10941373 PMID: [38491507](https://pubmed.ncbi.nlm.nih.gov/38491507/)

Trial design

900 participants in 3 patient groups

Standard MAP

Other group

Description:

Control group: MAP values between 50–60 mmHg

Treatment:

Other: Management of mean arterial pressure during cardiopulmonary by-pass in cardiac surgery interventions. The MAP will be maintained in the range of values of each study arm

High MAP

Other group

Description:

First Comparator group: MAP values between 70–80 mmHg

Treatment:

Other: Management of mean arterial pressure during cardiopulmonary by-pass in cardiac surgery interventions. The MAP will be maintained in the range of values of each study arm

Patient-tailored MAP

Other group


Description:

Second Comparator group: MAP comparable to the patient's pre-operative MAP. This one will be calculated by performing 3 blood pressure measurement in three different moments of the day before surgery (at 8 am, at 3 pm, and at 9 pm), and will be calculated using the standard formula "Diastolic AP + 0,33 x (systolic AP - Diastolic AP)". The preoperative MAP value obtained will be target during CPB, within a range of ± 10 mmHg

Treatment:

Other: Management of mean arterial pressure during cardiopulmonary by-pass in cardiac surgery interventions. The MAP will be maintained in the range of values of each study arm

Limites de la NIRS




Plus de désaturations cérébrales dans le groupe PAM haute


Impact of 2 Distinct Levels of Mean Arterial Pressure on Near-Infrared Spectroscopy During Cardiac Surgery: Secondary Outcome From a Randomized Clinical Trial

Frederik Holmgaard, BMSc,* Anne G. Vedel, MD,* Theis Lange, MSc, PhD,†‡
Jens C. Nilsson, MD, PhD,* and Hanne B. Ravn, MD, PhD, DMSc*

Cutaneous Vasoconstriction Affects Near-infrared Spectroscopy Determined Cerebral Oxygen Saturation during Administration of Norepinephrine

Henrik Sørensen, M.Sc.,* Niels H. Secher, M.D., D.M.Sc.,† Christoph Siebenmann, M.Sc.,‡
Henning B. Nielsen, M.D., D.M.Sc.,§ Matthias Kohl-Bareis, Ph.D.,|| Carsten Lundby, Ph.D.,#
Peter Rasmussen, Ph.D.**





Mort encéphalique: pas de perfusion sur la carotide interne
Clampage aortique:

Marqueur de perfusion tissulaire
Manque de spécificité de monitoring du tissu cérébral


Effect of Extracerebral Contamination on Near-infrared Spectroscopy as Revealed during Organ Donation: A Prospective Observational Study in Brain-dead Organ Donors

Martin Soehle, M.D., Ph.D., M.H.B.A.; Juliane Langer, M.S.; Ehrenfried Schindler, M.D., Ph.D.; Steffen Manekeller, M.D., Ph.D.; Mark Coburn, M.D., Ph.D.; Marcus Thudium, M.D., Ph.D.

+ Author and Article Information

Anesthesiology February 2024, Vol. 140, 231-239.

<https://doi.org/10.1097/ALN.0000000000004828>



Intégrer le débit de CEC dans la réflexion sur la PAM

- $PAM = DC * RVS * k$

Cerebral Oximetry and Mean Arterial Pressure: Not a Straight Relationship, the Flow Between?

Desebbe O, et al. Anesth Analg 2019. PMID 31743211

Circulation

ORIGINAL RESEARCH ARTICLE



High-Target Versus Low-Target Blood Pressure Management During Cardiopulmonary Bypass to Prevent Cerebral Injury in Cardiac Surgery Patients

A Randomized Controlled Trial

In 1928, Jarisch is quoted as saying, “It is a source of regret that the measurement of flow [*i.e.*, SV] is so much more difficult than the measurement of pressure. This has led to an undue interest in the blood pressure manometer. Most organs, however, require flow rather than pressure.

Table 2. Intraoperative Management

| | Low-Target Group (n=98) | High-Target Group (n=97) |
|---|-------------------------|--------------------------|
| Hematocrit, before start of surgery, % | 40.3±5.9 | 40.6±4.7 |
| MAP before anesthesia induction, mmHg | 92.3±15.7 | 96.9±13.4 |
| MAP during bypass, mmHg | 44.7±4.7 | 66.8±4.9 |
| MAP below target during bypass, n (%) [*] | 2 (2.0) | 18 (18.5) |
| MAP above target during bypass, n (%) [†] | 5 (5.1) | 0 (0) |
| Blood flow rate during bypass, L·min ⁻¹ ·m ⁻² | 2.69±0.1 | 2.69±0.1 |
| Hematocrit, mean level during bypass, % | 31.5±3.8 | 33.1±4.2 |
| Nadir hematocrit sampling value during bypass, % | 28.7±3.7 | 29.2±4.0 |
| Surgery time, min | 184.9±50.8 | 194.3±66.6 |
| Bypass time, min | 94.0±33.0 | 105.6±77.4 |
| Cross-clamp time, min [‡] | 63.3±26.9 | 64.8±32.6 |
| Peak lactate level during surgery, mmol | 2.25±0.83 | 2.16±0.82 |
| Norepinephrine infused in the OR, µg/kg | 2.65±6.01 | 17.43±20.14 |
| Patients receiving norepinephrine in the OR, n (%) | 35 (35.7) | 90 (92.7) |

**Improvement of Outcomes after
Coronary Artery Bypass II: A
Randomized Trial Comparing
Intraoperative High Versus Customized
Mean Arterial Pressure**

Mary E. Charlson, M.D.,* Janey C. Peterson, Ed.D., R.N.,* Karl H. Krieger, M.D.,†

Débits de CEC bas...

| B. Intraoperative Management Pre-bypass | High MAP | Custom MAP | P |
|--|-----------------|-------------------|----------|
| Pre-bypass MAP (mmHg) | 86 ± 8 | 85 ± 9 | |
| Pre-bypass cardiac output (L/min) | 4 ± 1 | 4 ± 1. | |
| Bypass Times (min) | | | |
| Bypass | 74 ± 24 | 77 ± 22 | ns |
| Cross-clamp | 40 ± 15 | 40.4 ± 14 | ns |
| Pump flows (L/min/m ²)# | | | |
| Bypass on—warming (± θ) | 2.0 ± 0.3 | 2.0 ± 0.3 | ns |
| Warming—cross-clamp off (± θ) | 2.1 ± 0.3 | 2.2 ± 0.3 | ns |
| Cross-clamp off—bypass off (± θ) | 2.4 ± 0.2 | 2.34 ± 0.3 | ns |

| | | n | population | Basse PAM | Haute PAM | Débit (l/min/m ²) | particularités |
|---|------|-----|---------------------------|-----------|-------------------------------------|---|--|
| Azau | 2014 | 300 | À risque d'AKI | 50-60 | 75-85 | 2,4 pour SvO ₂ > 70% 2,57 / 2,58 | Circuit clot, miniCEC  |
| Siepe | 2011 | 92 | Chir coronaire | 60-70 | 80-90 | 2,6 | Troubles cognitifs  |
| Charlson | 2007 | 412 | Chir coronaire programmée | 80 | +/- 20 mmHg, max 90 (57-90), 88 moy | 2 à 2,4  | 61% adhérence gp PAM préop 85% adhérence gp Pam 80 management at higher MAP on CPB dramatically reduces stroke occurrence in patients with severe aortic atheromatous disease (grade IV and V) |
| le débit de pompe est primordial | | | | | | | |
| Gold | 1995 | 248 | Chir coronaire | 50-60 | 80-100 | 1,9 à 2,3 | Niveau de PAM respectés  |
| Vedel | 2018 | 197 | Coronaires + valves | 40-50 | 70-80 | 2,7 |  |

Le débit de CEC avant la pression ?...

Observational Study




> J Cardiothorac Vasc Anesth, 32 (2), 684-690 Apr 2018

Disturbances in Oxygen Balance During Cardiopulmonary Bypass: A Risk Factor for Postoperative Delirium

Nina Smulter ¹, Helena Claesson Lingehall ², Yngve Gustafson ³, Birgitta Olofsson ⁴, Karl Gunnar Engström ⁵, Micael Appelblad ⁶, Staffan Svenmarker ⁶

Goal-directed perfusion to reduce acute kidney injury: A randomized trial

Marco Ranucci, MD, FESC,^a Ian Johnson, CCP,^{b,c} Timothy Willcox, CCP,^{d,e} Robert A. Baker, PhD, CCP,^f



Acute kidney injury in the goal-directed perfusion and control groups.

FIGURE 2. Oxygen delivery (DO_2) values in the goal-directed perfusion (blue; A) and control (red; B) arms during cardiopulmonary bypass (CPB). Boxes represent interquartile range, lines in the boxes represent the median, whiskers are 95% confidence intervals, and dots are outliers. The green line represents the critical DO_2 value of $280 \text{ mL} \cdot \text{min}^{-1} \cdot \text{m}^{-2}$. Data are restricted to the first 120 minutes of CPB.

Etude multicentrique,

Valeurs hautes de DaO2 dans les 2 bras.... Et pourtant moins d'Insuffisance rénale Aigue dans le groupe GDP

PAM dans les 2 groupes?

[The Effect of Pump Flow on Cerebral Oxygen Metabolism During Cardiopulmonary Bypass]

[Article in Japanese]

H Sakahashi¹

2.2-2.5 L/min/m² in group L and 2.7-3.0 L/min/m² in group H

It is suggested that cerebral metabolism should be decreased during cooling to 31 degrees C of pharyngeal temperature, 2.2-2.5 l/min/m² of pump flow was adequate to keep SjO₂ stable. On the other hand, it is necessary to increase pump flow to 2.7-3.0 l/min/m² during rewarming period as

**Le débit de CEC doit s'adapter à la demande métabolique
variabilité du débit**





Fig. 2 Cardiac index (CI), relative mean middle cerebral arterial velocity (%MCAV) and mean arterial pressure (MAP) : group L (●) and group H (■).

$$\%MCAV = MCAV \div MCAV_{pre-CPB} \times 100$$

Essayer de rester physiologique
= s'adapter aux besoins métaboliques



The Journal of Thoracic and Cardiovascular
Surgery

Volume 134, Issue 3, September 2007, Pages 587-593



Cardiopulmonary support and physiology

Elevated flow rate during cardiopulmonary bypass is associated with fluid accumulation

Oddbjørn Haugen MD ^a, Marit Farstad MD, PhD ^c, Venny Kvalheim MD ^b, Olav Bøe DDS, MSc ^d,
Paul Husby MD, PhD ^c  


Type de CEC supérieur à un autre pour préserver la PAM? Limiter les contraintes vasculaires par la miniCEC?

A combination of MiECC features, such as coating, centrifugal pump, separation of cardiotomy suction blood, and use of closed systems, should be considered to improve conventional CPB

Journal of Cardiothoracic and Vascular Anesthesia 35 (2021) 1–17



Fig. 1 Minimally invasive extracorporeal circulation (MiECC) during myocardial revascularization in end-stage coronary artery disease patients



Both ECC systems, MECC (minimized extracorporeal system) and CCPB (conventional cardiopulmonary bypass).

R4.4 – Il est recommandé de privilégier une technique de « CEC optimisée » afin de réduire la survenue de complications postopératoires et la mortalité hospitalière.

GRADE 1+ (accord FORT)

Minimal invasive extracorporeal circulation (MiECC): the state-of-the-art in perfusion

Kyriakos Anastasiadis, Helena Argiriadou, Apostolos Deliopoulos, Polychronis Antonitsis

Cardiothoracic Department, AHEPA University Hospital, Thessaloniki, Greece
 Correspondence to: Polychronis Antonitsis. Cardiothoracic Department, AHEPA University Hospital, S. Kyriakidi 1, 546 36 Thessaloniki, Greece.
 Email: antonits@auth.gr

Submitted Jan 08, 2019. Accepted for publication Jan 15, 2019.






Figure 1 Established classification of MiECC circuits (3). X, pump; O, oxygenator; C, cardioplegia; T, bubble-trap/air removing device; V, vent (aortic/pulmonary); S, soft-bag/reservoir; H, hard-shell/reservoir; MiECC, minimal invasive extracorporeal circulation.

miniCEC et PAM



If the mean arterial pressure decreased, the arterial flow was increased up to a maximum of 120% of the calculated flow. If mean arterial pressure further decreased, a 5 µg norepinephrine bolus was given to raise the MAP on CCPB between 45 and 75 mmHg.



miniCEC et PAM: causes d'une PAM plus élevée?



Article
Efficacy of Pulsatile Flow Perfusion in Adult Cardiac Surgery: Hemodynamic Energy and Vascular Reactivity

Mikhail Dodonov ¹, Francesco Onorati ¹, Giovanni Battista Luciani ¹, Alessandra Francica ^{1,*} , Maddalena Tessari ¹, Tiziano Menon ¹, Leonardo Gottin ², Aldo Domenico Milano ³ and Giuseppe Faggian ¹

- A new MEDOS Delta Stream DP3 centrifugal blood pump was used during CPB and was set in pulsatile (PP group) or non-pulsatile (NP group) mode.
- In pulsatile mode, the pump was set at maximum rotation speed variation of 3500–8500 rpm with a pulse frequency of 60 bpm; start and stop points of the pulse were set at 20 and 80% of the pulse cycle,
- This model includes the hemodynamic energy gradient as a key point rather than the pressure gradient and it is actually described in terms of energy equivalent pressure (EEP) and surplus hemodynamic energy (SHE). EEP was calculated according to the following formula: $EEP = Rtt_{12} f pdt / Rtt_{12} f dt$ (mm Hg), where f is blood flow, p is blood pressure and the product of flow and pressure represents hemodynamic power



| | | | |
|---|---------|-----------|------|
| Post-op CCr, 18 h (mL/min/1.73 m ²) | 61 ± 35 | 47 ± 16 * | 0.06 |
|---|---------|-----------|------|

| | | Start CPB | 1s of Cross-Clamp | 15s of Cross-Clamp | 30s of Cross-Clamp | Re-Warming | Cross-Clamp Removal | Means for all CPB Period | ANOVA p (Group/Time Interaction) |
|------------|----|-----------|-------------------|--------------------|--------------------|------------|---------------------|--------------------------|------------------------------------|
| MAP (mmHg) | PP | 57 ± 15 | 58 ± 14 | 51 ± 11 | 58 ± 12 | 58 ± 14 | 51 ± 11 | 56 ± 9.0 | 0.03 |
| | NP | 59 ± 12 | 61 ± 15 | 64 ± 19 ** | 69 ± 19 * | 61 ± 18 | 57 ± 12 | 62 ± 12 * | |


Intérêt d'une CEC pulsatile?



Contents lists available at [ScienceDirect](#)

Journal of Cardiothoracic and Vascular Anesthesia

journal homepage: www.jcvaonline.com



Editorial

Guidelines for Conduct of Cardiopulmonary Bypass

- Pulsatile perfusion may reduce postoperative pulmonary and renal complications and should be considered in patients at high risk for adverse lung and renal outcomes (grade B)



Autres voies pour optimiser la
PAM en chirurgie cardiaque?

Closed-Loop System of Vasopressor Infusion Post Cardiac Surgery

Currently, vasopressor infusions for managing hypotension post-surgery are adjusted manually



A novel closed-loop vasopressor (CLV) has been developed by physicians



Can the CLV replace manual titration?



Randomized trial of 42 patients after cardiac surgery



Norepinephrine infusion

Two-hour study period


CLV system
Manual (control)



Mean arterial pressure measured




% of time with hypotension (MAP <65 mmHg)



CLV - 1.4% | Control - 12.5%




% of time with MAP 65-75 mmHg



CLV - 95% | Control - 66%



% of time with MAP >75 mmHg



CLV - 3.2% | Control - 20.6%

Closed-loop vasopressor is an efficient system to control post-operative hypotension in patients with cardiac surgeries

WARNING sur les vasoconstricteurs?


CORRESPONDENCE

Open Access

Mean arterial pressure during cardiopulmonary bypass: A modifiable risk factor for acute kidney injury in cardiac surgery patients?

Nikolaus Schreiber¹, Simon Orlob¹, Stephanie Fida¹, Christoph Klivinyi¹, Alexander H. Kirsch², Michael Kolland² and Michael Schörghuber^{1*}

b. Predicted probabilities for development of CSA-AKI with increasing TWA-MAP < 65mmHg during CPB




Besides severity of hypotension, we observed a significant association between increased vasopressor dosage and CSA-AKI development in multivariable analysis, which may reflect the intricate relationship between renal perfusion, vasopressor exposition and consecutive CSA-AKI, meriting further research.

WARNING sur les vasoconstricteurs?


ANESTHESIOLOGY

Intraoperative Hypotension and Acute Kidney Injury, Stroke, and Mortality during and outside Cardiopulmonary Bypass: A Retrospective Observational Cohort Study

Miguel Armengol de la Hoz, M.S.,
Valluvan Rangasamy, M.D., D.N.B., D.E.S.A.,
Andres Brenes Bastos, M.D., Xinling Xu, Ph.D.,
Victor Novack, M.D., Bernd Saugel, M.D.,
Balachundhar Subramaniam, M.D., M.P.H., F.A.S.A.
ANESTHESIOLOGY 2022; 136:927–39



WARNING sur les vasoconstricteurs?



3. Postoperative complications according to norepinephrine use in the propensity score-matched cohort. Major adverse kidney and cerebral events include stroke, acute kidney injury, and death.



Original Article

Epidemiology, risk factors and outcomes of norepinephrine use in cardiac surgery with cardiopulmonary bypass: a multicentric prospective study

Pierre-Grégoire Guinot^{a,b,*}, Bastien Durand^a, Emmanuel Besnier^c, Paul-Michel Mertes^d, Chloe Bernard^e, Maxime Nguyen^{a,b}, Vivien Berthoud^a, Osama Abou-Arab^f, Belaid Bouhemad^{a,b}, Collaborator study group¹



2500 patients,
étude multicentrique
observationnelle prospective,
Utilisation score de propensité

Travailler sur la PAM centrale...?

Open access

Protocol

BMJ Open Femoral versus radial arterial pressure monitoring in cardiac surgery patients: protocol for a randomised controlled multicentric superiority trial (FERARI)

Janvier 2026
2 centres français

Pierre-Grégoire Guinot,^{1,2,3} Romain Bronnert,^{4,5,6} Tommy Grelet,⁷
Belaid Bouhemad,^{1,2,3} Maxime Nguyen,^{1,2,3} Guillaume Besch ,^{4,5,6} FERARI study group

Objective To determine whether femoral arterial pressure monitoring reduces norepinephrine use compared with radial monitoring in cardiac surgery patients.



Les nouveaux médicaments vasoconstricteurs

Table 2.

Therapeutic options for management of vasoplegia after cardiac surgery.^{1,10,42} NO: nitric oxide; V1: vasopressin; AT1: angiotensin 1

| TREATMENT OPTION | RECOMMENDED DOSAGE | MECHANISM OF ACTION |
|---------------------------------|---|--|
| Catecholamines | | |
| Norepinephrine | 0.01-0.2 µg/kg/min | Strong α-1, α-2, and mild β-1 adrenergic agonist |
| Epinephrine | 0.01-0.5 mcg/kg/min | Strong α-1, α-2, and β-1 adrenergic agonist |
| Dopamine | 0-20 mcg/kg/min | α-1 adrenergic agonist at higher doses (dose-dependent agonism) |
| Phenylephrine | 0.5-5 µg/kg/min | α-1 and α-2 adrenergic agonist |
| Non-catecholamine agents | | |
| Vasopressin | 0.03 U/min | Acts on <i>AVPR1a</i> receptor to reduce vasodilatory effects of NO; V1 agonist |
| Methylene blue | 1.5-2.0 mg/kg/bolus over 15 to 30 minutes | Inhibits guanylyl cyclase and endothelial NO synthase |
| Angiotensin 2 | Start 20 ng/kg/min | AT1 agonist; stimulates release of vasopressin from posterior pituitary gland and release of aldosterone from adrenal gland |
| Hydroxocobalamin | 5 g infusion over 5 min | Inhibits guanylyl cyclase, endothelial NO synthase, and hydrogen sulfide |
| Ascorbic acid | 6 g IV bolus/day | Acts as a cofactor for catecholamine synthesis |
| Hydrocortisone | 50 mg bolus every 6 h or 100 mg every 8 h (10 mg/h) | Hydrocortisone in addition to vitamin C, and thiamine decrease required dose for vasopressors; improves catecholamine response |

Angiotensine II

- **PORTHOS Trial – ACTRN12623000848606**
 - Design: essai randomisé, en double aveugle, multicentrique, comparant une perfusion d'angiotensine II versus noradrénaline, en per-op et jusqu'à 48 h après le début de la chirurgie cardiaque sous CEC. [pmc.ncbi.nlm.nih](#)
 - Population: 400 patients de chirurgie cardiaque à haut risque d'AKI (critères: Hb <130 g/L, créatinine >100 µmol/L, âge >70 ans, NYHA IV, IMC >30, au moins un critère). [pmc.ncbi.nlm.nih](#)
 - Objectif: maintenir une PAM 70–80 mmHg; critère principal = durée de séjour hospitalier, critères secondaires = événements rénaux, cardiovasculaires et neurologiques. [pmc.ncbi.nlm.nih](#)
 - Statut: protocole publié, essai en cours (pas encore de résultats cliniques définitifs). [pmc.ncbi.nlm.nih](#)

Vasopressine

Randomized Controlled Trial > Am Heart J. 2024 Jun;272:86-95. doi: 10.1016/j.ahj.2024.03.008.

Epub 2024 Mar 16.

Prospective randomized double-blind study to evaluate the superiority of Vasopressin versus Norepinephrine in the management of the patient at renal risk undergoing cardiac surgery with cardiopulmonary bypass (NOVACC trial)

Pierre-Grégoire Guinot ¹, Olivier Desebbe ², Guillaume Besch ³, Philippe Guerci ⁴,
Philippe Gaudard ⁵, Diane Lena ⁶, Paul Michel Mertes ⁷, Osama Abou-Arab ⁸, Belaid Bouhemad ⁹;
NOVACC study group

Vp mieux pour le rein?: PMID: **41208542**

conclusion



RECOMMANDATIONS FORMALISEES D'EXPERTS

Réhabilitation Améliorée Après Chirurgie Cardiaque adulte sous CEC ou à cœur battant

ENHANCED RECOVERY AFTER CARDIAC SURGERY UNDER CPB OR OFF-PUMP

2021

RFE commune SFAR - SFCTCV
Société Française d'Anesthésie et de Réanimation
Société Française de Chirurgie Thoracique et Cardiovasculaire

Special Communication

FREE

Cite Permissions Metrics

Guidelines for Perioperative Care in Cardiac Surgery Enhanced Recovery After Surgery Society Recommendations

JAMA Surg
Published Online: May 4, 2019
2019;154;(8):755-766. doi:10.1001/
jamasurg.2019.1153

Daniel T. Engelman, MD¹; Walid Ben Ali, MD²; Judson B. Williams, MD, MHS³; et al

» Author Affiliations | Article Information

Pas de recommandations sur la PAM perCEC ou la DaO2

CHAMP 4. STRATEGIE CHIRURGICALE ET GESTION DE LA CEC

| | | | |
|-------|---|--------------|------|
| 4.1 | La chirurgie mitrale vidéo-assistée peut être envisagée dans des équipes entraînées | Avis experts | FORT |
| 4.2 | Réaliser la CEC en normothermie | 1+ | FORT |
| 4.3.1 | Ne pas réaliser systématiquement les pontages coronariens à cœur battant | 1- | FORT |
| 4.3.2 | Les pontages coronariens à cœur battant peuvent être discutés en cas d'aorte très calcifiée | Avis experts | FORT |
| 4.4 | Utiliser une CEC optimisée | 1+ | FORT |
| 4.5 | Ne pas privilégier une technique de cardioplégie plutôt qu'une autre | 2- | FORT |

► Interdiscip Cardiovasc Thorac Surg. 2025 Feb 14;40(2):ivaf002. doi: [10.1093/icvts/ivaf002](https://doi.org/10.1093/icvts/ivaf002) 

2024 EACTS/EACTAIC/EBCP Guidelines on cardiopulmonary bypass in adult cardiac surgery

In summary, a high blood pressure target may result in little to no difference in patient outcomes, including AKI and deaths and further studies are needed to assess the efficacy of a higher blood pressure target among those who undergo cardiac surgery with CPB. It is more important to target DO_2 , optimize blood flow from CPB and monitor markers of tissue perfusion (NIRS and $SvcO_2$) rather than use vasopressors to increase ABP (please see also section 9.7 on goal-directed perfusion).

Conclusion

- PAM: un outil de surveillance parmi d'autre: PaCO₂, débit CEC, SvO₂, Hte
- La PAM est la résultante du Débit de CEC et des RVS :
 - PAM minimale si débit de CEC suffisant +++
 - Coupler la PAM à la DaO₂ +++ et à la demande métabolique (test de PAM sur la NIRS, sur la SVO₂, pas sur l'exCO₂...)
- Modifier les RVS dans un deuxième temps seulement
- Quel vasopresseur idéal?
- Titrer la PAM
 - sur organe « cible », sur microcirculation
 - maintenir PAM > 50 voire 65 mmHg sinon
 - Difficile de proposer une valeur cible de PAM selon PAM préopératoire

Proposition personnelle d'algorithme, en normothermie

Hématocrite >
18-24%

| Recommendations | Class ^a | Level ^b | Ref ^c |
|--|--------------------|--------------------|------------------|
| PRBC transfusions | | | |
| It is recommended that PRBCs be transfused during CPB if the Hb value is <6.0 g/dl. | I | C | |
| For HCT values between 18% and 24%, PRBCs may be transfused based on an assessment of the adequacy of tissue oxygenation. ^d | IIb | B | [243] |
| PRBCs should not be transfused during CPB if the HCT is >24%. | III | C | |

FiO2 pour SpO2 < 99%
(ne pas surestimer la SvO2)

Débit de CEC
pour SvO2 >
75%

Perfusion adaptée et non
luxuriante si réflexion sur
niveau de DaO2 seul

Débit de CEC
pour SvO2 >
75%



PAM
40-80 mmHg



1 PaCO2
40-45 mmHg
2 PVC < 10
3 SpO2 < 98%
4 Sédation adapté (40-60 pas de BSR)

NIRS > 80%
valeur de
base?

oui

non

DaO2 > 300
ml/min/m²

Test
noradrénaline

