université
de **BORDEAUX**

Sevrage respiratoire DU Kiné

Mai 2024

Pr Hadrien Rozé
Réanimation Polyvalente,
Centre Hospitalier Côte Basque
Université de Bordeaux

LES GRANDS PRINCIPES DU SEVRAGE

- ***Simple à Difficile***
- ***Sédation***
- ***Fonction Cardiaque***
- ***Force musculaire***

Weaning from mechanical ventilation in intensive care units across 50 countries (WEAN SAFE): a multicentre, prospective, observational cohort study

Tài Pham, Leo Heunks, Giacomo Dellani, Fabiana Madotto, Irene Aragao, Gaetan Deduneau, Ewan C Goligher, Giacomo Grasselli, Jon I Henrik Laake, Jordi Manóvilha, Oscar Peñaletas, Lise Piquillaud, Antonia Presenti, Hannah Wunsch, Frank van Haren, Laurent Brochard*, John Gaffney*, for the WEAN SAFE Investigators†

Lancet Respir Med 2023;
11: 465-76

Sevrage 3 situations

1) **Sevrage simple** (premier essai)

65%

2) **Sevrage prolongé** (plus d'un essai)

25%

3) **Sevrage (très) difficile**

10%

Sevrage 3 situations

	Funk	Penuelas	Sellares	Tunnelier
Simple	59 %	55 %	44 %	30 %
Prolongé	26 %	39 %	37 %	40 %
Difficile	14 %	6 %	18 %	30 %

Mécanismes du Sevrage non difficile

- Sevrage possible d'emblée:
 - * screening quotidien
- Sevrage possible mais non identifié
 - * Sur assistance
 - * Trop de sédation
 - * Pas de screening

Mécanismes du Sevrage difficile

- Pathologie pulmonaire/cardiaque sous jacente sévère
- Surcharge volémique
- Faiblesse musculaire respiratoire
- Métabolique/Nutrition/Anémie

TASK FORCE

Weaning from mechanical ventilation

J.-M. Boles^{*}, J. Bion[†], A. Connors[‡], M. Herridge[§], B. Marsh[¶], C. Melot^{||}, R. Pearl^{**},
H. Silverman^{##}, M. Stanchina^{††}, A. Vieillard-Baron⁺⁺, T. Welte^{§§}

TABLE 3 Classification of patients according to the weaning process

Group/category	Definition
Simple weaning	Patients who proceed from initiation of weaning to successful extubation on the first attempt without difficulty
Difficult weaning	Patients who fail initial weaning and require up to three SBT or as long as 7 days from the first SBT to achieve successful weaning
Prolonged weaning	Patients who fail at least three weaning attempts or require >7 days of weaning after the first SBT

SBT: spontaneous breathing trial.

Commande centrale
Amplitude, timing, distribution

Tronc cérébrale

Transmission nerveuse périphérique
Propagation du potentiel d'action
Jonction neuro-musculaire
Excitation fibre musculaire

Activation électrique musculaire
Propriétés contractiles
(relation longueur-tension, fatigue)

Contraction musculaire (force)
Configuration de la cage thoracique
Couplage neuro mécanique

Distension pulmonaire
Pression pleurale diminuée
Auto PEEP
Elastance thoraco pulmonaire (P_{totE})

Diminution Palvéolaire
Résistances voies aériennes (P_{totR})

Pression ouverture voies
aériennes, débit, volume
 VD/VT FR

Ventilation alvéolaire
Echanges gazeux, hématoxé

Protocole de sedation

Kress, JP. et al. N Engl J Med 2000; 342:1471-1477

Efficacy and safety of a paired sedation and ventilator weaning protocol for mechanically ventilated patients in intensive care (Awakening and Breathing Controlled trial): a randomised controlled trial

Lancet 2008; 371: 126–34

A protocol of no sedation for critically ill patients receiving mechanical ventilation: a randomised trial

Thomas Strøm, Torben Martinussen, Palle Toft

Lancet 2010

Test à l'anexate démontre l'inhibition de la sedation

REVIEW

Weaning failure of cardiac origin: recent advances

Jean-Louis Teboul¹, Xavier Monnet, and Christian Richard

Christian Richard
Jean-Louis Teboul

Weaning failure from cardiovascular origin

Invasive Crit Care Med
DOI 10.1007/s00134-019-1870-0

ORIGINAL

Belén Cabello
Arnand W. Thille
Ferran Rochecampo
Laurent Brochard
Francisco J. Gómez
Jordi Manóvilha

Physiological comparison of three spontaneous breathing trials in difficult-to-wean patients

Fonction cardiaque

Balance hydrique

Intensive Care Med (2005) 31:1643–1647
DOI 10.1007/s00134-005-2801-3

ORIGINAL

Anupama Upadya
Lisa Tilluckdharry
Visvanathan Muralidharan
Yaw Annateng-Adjepong
Constantine A. Manthous

Fluid balance and weaning outcomes

	Weaning success (n=39)	Weaning failure (n=48)	All patients (n=87)	<i>p</i>
Duration of ventilation (days)	2 (1 to 8)	3 (1 to 14)	3 (1 to 14)	0.03
APACHE II (day of BT)	17 (3 to 30)	15 (3 to 25)	16 (3 to 30)	0.2
Fluid balance 24 h (ml)	-625 (-4,380 to +3,274)	+242 (-3,923 to +4,272)	-91 (-4,380 to +4,272)	0.01
Cumulative fluid balance (ml)	-633 (-8,232 to +9,534)	+920 (-1,1760 to +2,0483)	-65 (-1,1760 to +2,0483)	0.06
Diuretics on day of trial	62%	51%	56%	0.2
Prealbumin (mg/dl)	14 (2 to 35)	12 (2 to 30)	13 (2 to 35)	0.8
f/V _t (breaths min ⁻¹ l ⁻¹)	50 (13 to 260)	80 (17 to 300)	67 (13 to 300)	0.005
PaO ₂ /FIO ₂	240 (88 to 477)	244 (100 to 503)	240 (88 to 503)	0.9
Compliance (ml/cmH ₂ O)	44 (22 to 76)	39 (13 to 80)	43 (13 to 80)	0.2
Left ventricular dysfunction (%)	69	68	69	0.6

Faiblesse musculaire: Dysfonction diaphragmatique

J Appl Physiol 92: 2585–2595, 2002.

N Engl J Med 2008;358:1327-35.

Rapidly Progressive Diaphragmatic Weakness and Injury during Mechanical Ventilation in Humans

Samir Jaber^{1,2,6}, Basl J. Petrof³, Boris Jung^{1,2}, Gérald Chanques^{1,2}, Jean-Philippe Berthet⁴, Christophe Rabuel⁵, Hassan Bouyabrine⁶, Patricia Courouble^{1,2}, Christelle Koechlin-Ramonatxo⁷, Mustapha Sebbane^{1,2}, Thomas Similowski⁸, Valérie Scheuermann⁹, Alexandre Mebazaa⁵, Xavier Capdevila^{1,2}, Dominique Mornet², Jacques Mercier^{4,10}, Alain Lacampagne⁹, Alexandre Philips², and Stefan Matecki^{2,10}

Respiratory weakness is associated with limb weakness and delayed weaning in critical illness*

Bernard De Jonghe, MD; Sylvie Bastuji-Garin, MD, PhD; Marie-Christine Durand, MD; Isabelle Malissin, MD; Pablo Rodrigues, MD; Charles Cerf, MD; Hervé Outin, MD; Tarek Sharshar, MD, PhD; for Groupe de Réflexion et d'Etude des Neuromyopathies En Réanimation

Table 6. Analysis of risk factors for low maximal inspiratory pressure (MIP) or maximal expiratory pressure (MEP)

Univariate Analysis	MIP or MEP $\leq$ 30 cm H ₂ O (n = 53)	MIP and MEP >30 cm H ₂ O (n = 26)	<i>p</i> Value
At ICU admission			
Age in years, median (IQR)	67 (53–78)	70 (54–76)	.9
Female sex, n (%)	22 (41.5)	7 (26.9)	.3
Admission SAPS II, median (IQR)	46 (36–58)	44 (38–50)	.4
COPD, n (%)	20 (37.7)	13 (50)	.3
Before awakening			
Days of MV, median (IQR)	10 (8–14)	10 (8–13)	.9
Days with $\geq$ 2 failed organs, median (IQR)	6 (4–10)	5 (0–9)	.2
Use of corticosteroids, n (%)	33 (62.3)	16 (61.5)	.9
Use of neuromuscular blockers, n (%) ^a	13 (24.5)	11 (42.3)	.1
Average daily morning BGL (mmol/L), median (IQR)	8.1 (7.0–9.3)	7.5 (6.8–8.6)	.1
Septic shock, n (%)	31 (58.5)	8 (30.8)	.02
Multivariate Logistic Regression Analysis of Low MIP or MEP $\leq$30 cm H₂O			
	OR	95% CI	<i>p</i> Value
Septic shock	3.17	1.17–8.58	.02

Table 5. Independent determinants of the risk of successful extubation delayed for >=7 days after awakening

	Model with MIP			Model with MEP			Model with VC			Model with MRC					
	OR	95% CI	P Value	OR	95% CI	P Value	OR	95% CI	P Value	OR	95% CI	P Value			
MIP <30 cm H ₂ O	8.02	2.12–30.36	.002	MEP <30 cm H ₂ O	4.15	1.16–14.82	.03	VC <11.1 mL/kg	2.75	0.82–9.18	.1	MRC <41	3.03	1.23–7.43	.02
COPD	4.43	1.20–16.41	.03	COPD	4.56	1.24–16.75	.02	COPD	4.43	1.3–14.79	.02	COPD	2.74	1.10–6.85	.03
Cardiac insufficiency	4.96	1.25–19.71	.02	Cardiac insufficiency	3.79	1.07–13.39	.04	Cardiac insufficiency	3.24	0.92–11.38	.07	Cardiac insufficiency	2.14	0.82–5.61	.1

RESEARCH

Open Access

Accuracy of respiratory muscle assessments to predict weaning outcomes: a systematic review and comparative meta-analysis

Diego Poddighe^{1,2†}, Marne Van Hullebeke^{1,2†}, Yasir Qaiser Choudhary¹, Debora Ribeiro Campos³, Michele R. Schaeffer¹, Jan Y. Verbakel^{4,5}, Greet Hermans^{7,6}, Rik Gosselink^{1,2,7} and Daniel Langer^{1,7*}

Maximal inspiratory pressure

Airway occlusion pressure

Diaphragm excursion

Study	TP	FP	FN	TN	n	P _{max} Threshold	CONDITION	Risk of bias	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Elgazzar 2019	16	7	3	4	30	9.1	SBT	Low	0.64 [0.60, 0.97]	0.96 [0.11, 0.99]		
Huang 2025	40	15	10	25	88	9.05	MV	High	0.60 [0.56, 0.90]	0.58 [0.41, 0.74]		
Kim 2011	11	0	17	45	82	10.0	SBT	Low	0.59 [0.22, 0.59]	0.83 [0.71, 0.92]		
Haji 2018	24	5	18	6	53	10.0	SRT	High	0.57 [0.41, 0.75]	0.55 [0.25, 0.83]		
Vivier 2019	105	13	50	14	189	10.0	SBT	High	0.68 [0.50, 0.75]	0.42 [0.25, 0.61]		
Raneek 2016	16	5	7	1	30	10.0	SRT	Low	0.70 [0.47, 0.80]	0.74 [0.60, 0.88]		
Vannr-Vega 2021	67	1	16	1	84	10.0	SHI	Low	0.81 [0.71, 0.89]	1.00 [0.83, 1.00]		
Osman 2017	42	3	9	18	69	10.0	SBT	Low	0.82 [0.69, 0.96]	1.00 [0.81, 1.00]		
Hevan 2016	17	1	3	6	27	10.0	SBT	Low	0.85 [0.62, 0.97]	0.85 [0.42, 1.00]		
Spadaro 2021	25	3	4	2	62	10.0	MV	Low	0.87 [0.69, 0.96]	0.05 [0.01, 0.21]		
Pelkar 2018	49	14	4	6	73	10.0	SBT	Low	0.92 [0.82, 0.96]	0.30 [0.12, 0.54]		
Elmadh 2019	16	2	1	11	30	10.0	SBT	Low	0.94 [0.71, 1.00]	0.65 [0.50, 0.98]		
Ekombetchal 2023	95	4	26	4	130	10.5	NR	High	0.78 [0.70, 0.86]	0.50 [0.10, 0.84]		
Huang 2017	10	7	2	21	40	10.7	SBT	Low	0.83 [0.52, 0.96]	0.75 [0.50, 0.89]		
Mariani 2015	21	0	11	2	34	11.0	SBT	Low	0.66 [0.47, 0.81]	1.00 [0.16, 1.00]		
Keur 2022	20	2	5	17	50	11.0	GGT	High	0.64 [0.50, 0.90]	0.09 [0.07, 0.89]		
Jiang 2004	27	4	5	19	55	11.0	SBT	High	0.64 [0.67, 0.95]	0.63 [0.61, 0.95]		
Seeed 2010	19	1	3	7	30	11.0	SBT	High	0.66 [0.55, 0.97]	0.60 [0.47, 1.00]		
Ferghay 2017	35	4	5	10	54	11.0	SBT	High	0.68 [0.70, 0.96]	0.71 [0.42, 0.92]		
Benejee 2018	38	3	2	10	63	11.0	SBT	Low	0.68 [0.83, 0.96]	0.77 [0.46, 0.96]		
Holmy 2021	0	1	0	12	22	11.0	SBT	Low	0.68 [0.68, 1.00]	0.02 [0.04, 1.00]		
Xu 2022	33	3	26	26	69	11.0	SBT	High	0.71 [0.62, 0.80]	0.79 [0.62, 0.90]		
Alam 2022	14	2	4	11	31	11.0	SBT	Low	0.72 [0.52, 0.94]	0.85 [0.64, 0.98]		
Samana 2017	24	14	9	17	64	11.0	SBT	High	0.73 [0.54, 0.87]	0.55 [0.36, 0.73]		
Hivoli 2023	25	4	9	6	43	12.0	SRT	High	0.74 [0.56, 0.87]	0.58 [0.21, 0.86]		
Hoyal 2017	60	7	16	17	100	12.0	SHI	High	0.78 [0.68, 0.87]	0.71 [0.49, 0.87]		
Shamil 2022	36	0	5	6	53	12.0	SHI	High	0.88 [0.74, 0.96]	1.00 [0.66, 1.00]		
Fossat 2022	75	15	7	1	100	12.0	SBT	High	0.92 [0.83, 0.97]	0.05 [0.00, 0.29]		
Arbas 2018	34	1	3	12	50	12.0	SBT	Low	0.92 [0.78, 0.98]	0.92 [0.64, 1.00]		
Sarawan 2022	160	3	11	21	200	12.1	SBT	High	0.94 [0.89, 0.97]	0.72 [0.50, 0.87]		
Vieira 2022	33	14	12	22	81	12.5	SBT	High	0.73 [0.58, 0.80]	0.61 [0.43, 0.77]		
ElsFady 2020	33	0	1	20	62	12.5	SBT	Low	0.97 [0.85, 1.00]	0.62 [0.60, 0.94]		
Luo 2017	34	2	16	6	60	12.6	SBT	High	0.68 [0.53, 0.80]	0.80 [0.44, 0.97]		
Therawit 2016	25	2	23	8	62	12.8	SBT	Low	0.55 [0.40, 0.66]	0.62 [0.46, 0.86]		
Mawla 2022	31	0	20	36	90	13.0	SBT	Low	0.61 [0.46, 0.74]	1.00 [0.91, 1.00]		
Li 2021	45	12	24	26	101	13.0	GGT	Low	0.65 [0.53, 0.76]	0.63 [0.44, 0.79]		
Gok 2021	20	2	12	4	40	13.1	GGT	High	0.70 [0.53, 0.80]	0.07 [0.22, 0.90]		
Scng 2022	65	13	8	24	110	13.5	GGT	High	0.89 [0.80, 0.90]	0.65 [0.47, 0.80]		
Spadaro 2016	21	2	10	10	51	14.0	BBT	High	0.62 [0.44, 0.70]	0.00 [0.04, 0.89]		
Yeo 2018	31	2	16	11	60	14.0	SBT	High	0.66 [0.51, 0.70]	0.85 [0.65, 0.98]		
Khan 2018	46	7	16	21	90	14.0	BBT	High	0.74 [0.62, 0.84]	0.75 [0.55, 0.89]		
Thabet 2020	14	0	7	10	40	14.1	SBT	Low	0.67 [0.43, 0.86]	0.63 [0.20, 0.76]		
Zayroun 2021	8	1	29	37	61	16.0	SBT	High	0.23 [0.10, 0.30]	0.06 [0.02, 1.00]		
Ali 2017	25	4	3	22	64	16.0	MV	High	0.86 [0.72, 0.96]	0.85 [0.65, 0.96]		
Al Taysr 2022	11	2	5	6	24	16.0	SBT	High	0.60 [0.41, 0.80]	0.75 [0.35, 0.97]		
Seeed 2010	23	1	1	7	32	16.0	MV	Low	0.96 [0.79, 1.00]	0.89 [0.47, 1.00]		
Mohamed 2021	41	7	19	13	80	17.0	SRT	High	0.68 [0.55, 0.80]	0.65 [0.41, 0.85]		
Zhang 2020	19	3	6	6	37	17.2	SHI	High	0.76 [0.55, 0.91]	0.75 [0.45, 0.95]		
Amara 2022	11	21	5	44	81	17.9	NR	Low	0.69 [0.41, 0.85]	0.88 [0.55, 0.79]		
Abdelhakez 2019	117	5	0	116	240	18.4	SBT	High	1.00 [0.97, 1.00]	0.96 [0.91, 0.98]		
El 2022	13	2	6	6	27	22.1	SBT	High	0.68 [0.43, 0.84]	0.75 [0.30, 0.94]		
Carrie 2017	32	3	13	13	61	27.0	SBT	Low	0.71 [0.56, 0.84]	0.99 [0.36, 0.79]		
Sead 2022	7	3	5	40	60	61.0	SBT	Low	0.58 [0.28, 0.80]	0.83 [0.70, 0.93]		

DTF n°1

Faiblesse musculaire

Review

J Appl Physiol 107: 962–970, 2009.
First published April 30, 2009; doi:10.1152/jappphysiol.00165.2009.

HIGHLIGHTED TOPIC | *The Respiratory Muscles in Chronic Obstructive Pulmonary Disease*

Role of the respiratory muscles in acute respiratory failure of COPD:
lessons from weaning failure

Martin J. Tobin,¹ Franco Laghi,¹ and Laurent Brochard²

Is Weaning Failure Caused by Low-Frequency Fatigue of the Diaphragm? Voir de la fatigue ?

Franco Laghi, Steven E. Cattapan, Amal Jubran, Salram Parthasarathy, Paul Warshawsky, Yoon-Sub A. Chol, and Martin J. Tobin

Am J Respir Crit Care Med Vol 167. pp 120–127, 2003

Inspiratory Muscle Rehabilitation in Critically Ill Adults

A Systematic Review and Meta-Analysis

Stefannie Vorona¹, Umberto Sabatini¹, Sulalman Al-Maqball¹, Michele Berton¹, Martin Dres^{2,3}, Bernie Blissett^{1,5}, Frank Van Haren^{5,6,7}, A. Daniel Martin⁸, Cristian Urrea¹, Debbie Brace¹, Matteo Parotto^{9,10,11}, Margaret S. Hemidge^{1,9,12}, Neill K. J. Adhikari^{9,13,14}, Eddy Fan^{1,9,12,15}, Luana T. Melo¹⁶, W. Darlene Reid^{9,16}, Laurent J. Brochard^{2,9,12}, Niall D. Ferguson^{1,9,12,14,15}, and Ewan C. Goligher^{1,9,15}

Outcome	Impact Effect (95% CI)	No. of Participants (RCTs)	Quality of the Evidence (GRADE)
Change in maximal inspiratory pressure from baseline after IMT	Mean difference in change 6 (5 to 8) cm H ₂ O higher in IMT group than in control group Pooled ratio of means for change in MIP relative to baseline MIP, 1.21 (1.16 to 1.26)	647 (15 RCTs)	⊗○○○ Very low ^{++‡}
Change in maximal inspiratory pressure from baseline after IMT (sensitivity analysis excluding studies at high risk of bias)	Mean difference 9 (7 to 12) cm H ₂ O higher in IMT group than in control group	175 (3 RCTs)	⊕⊕⊕⊕ High
Maximal inspiratory pressure after IMT	Mean difference 7 (6 to 8) cm H ₂ O higher in IMT group than in control group	575 (15 RCTs)	⊕⊕○○ Low ⁺⁺
Change in maximal expiratory pressure from baseline after IMT	Mean difference in change 9 (5 to 14) cm H ₂ O higher in IMT group than in control group Pooled ratio of means for change in MEP relative to baseline MEP, 1.39 (1.27 to 1.54)	153 (4 RCTs)	⊕⊕⊕○ Moderate [*]
Change in maximal expiratory pressure from baseline after IMT (sensitivity analysis excluding studies at high risk of bias)	Mean difference in change 9 (5 to 14) cm H ₂ O higher in IMT group than in control group	106 (2 RCTs)	⊕⊕⊕⊕ High
Duration of ventilation	Pooled duration of ventilation was 4.1 (0.8 to 7.4) d shorter in IMT group than in control group	325 (9 RCTs)	⊗○○○ Very low ^{++‡§}
Duration of ventilation (sensitivity analysis excluding studies at high risk of bias)	Pooled duration of ventilation was 4.6 (-1.0 to 10.1) d shorter in IMT group than in control group	220 (4 RCTs)	⊕⊕○○ Low ^{†§}
Duration of weaning from mechanical ventilation	Pooled duration of weaning from mechanical ventilation was 2.3 (0.7 to 3.9) d shorter in IMT group than in control group	257 (8 RCTs)	⊗○○○ Very low ^{++§}
Duration of weaning (sensitivity analysis excluding studies at high risk of bias)	Pooled duration of weaning from mechanical ventilation was 3.2 (0.6 to 5.8) d shorter in IMT group than in control group	208 (5 RCTs)	⊕⊕○○ Low ^{†§}
ICU length of stay	Length of stay in ICU was 3.1 (-1.0 to 7.1) d shorter in IMT group than in control group	28 (2 RCTs)	⊗○○○ Very low ^{++§}
Mortality in ICU	Pooled relative risk of death in ICU was 0.67 (0.20 to 2.20) in IMT group compared with control group	197 (3 RCTs)	⊕⊕○○ Low ^{†§}

The impact of inspiratory muscle training (IMT) on the duration of weaning from mechanical ventilation

ur

(1) Recherche quotidienne du pré-requis à l'épreuve de ventilation spontanée par le personnel soignant

- Absence d'inotrope et de vasopresseur
- Absence de sédation
- Réponse cohérente aux ordres simples
- $FiO_2 < 50\%$
- $PEP < 5\text{ cmH}_2\text{O}$

PEEP < 10

**(2) Épreuve de ventilation spontanée (VS) :
pièce en T au Aide Inspiratoire (AI) sans PEP ($6 \leq AI \leq 8\text{ cmH}_2\text{O}$)
+ recherche de signes de mauvaise tolérance**

- $FR > 35/\text{min}$
- $SpO_2 < 90\%$
- Variation de plus de 20 % de FC ou PAS
- Sueurs, agitation, troubles de la vigilance

Présents
Gaz du sang
et reprise de la VM

(3) Absents
Recherche des
critères d'extubation

**(4) Sevrage
difficile**

Extubation

> 30 j

48 h

**(6) Échec
du sevrage**

**(5) Sevrage
réussi**

Réintubation
ou VNI

LA VENTILATION ASSISTEE LORS DU SEVRAGE

- ***Niveau d'aide***
- ***Synchronisation***

Problématiques du mode lors du sevrage

- Synchroniser le patient au respirateur/limiter les asynchronies
- Adapter l'aide inspiratoire aux besoins du patient
 - VSAI: aide fixe à adapter manuellement
 - Intelligence artificielle en VSAI: Smartcare
 - Neurally Adjusted Ventilatory Assist: NAVA
 - Proportionnal Assist Ventilation: PAV+

N Engl J Med 2013;369:2126-36

JAP 2002;92:2585-2595

Dysfonctions

Niveau d'assistance

Intensive Care Med 2006;32:1515-1522

NARRATIVE REVIEW

Clinical strategies for implementing lung and diaphragm-protective ventilation: avoiding insufficient and excessive effort

$$P_{mus} + P_{vent} = \text{Volume} \times E + \text{Débit} \times R$$

PAV+ et NAVA (P_{aw})

VSAI (P_{aw})

Effort (P_{mus})

The New England Journal of Medicine

©Copyright, 1995, by the Massachusetts Medical Society

Volume 332

FEBRUARY 9, 1995

Number 6

A COMPARISON OF FOUR METHODS OF WEANING PATIENTS FROM MECHANICAL VENTILATION

ANDRÉS ESTEBAN, M.D., Ph.D., FERNANDO FRUTOS, M.D., MARTÍN J. TORRES, M.D., INMACULADA ALÍA, M.D., JOSÉ F. SOLSONA, M.D., INMACULADA VALVERDÚ, M.D., RAFAEL FERNÁNDEZ, M.D., MIGUEL A. DE LA CAL, M.D., SALVADOR BENITO, M.D., Ph.D., ROGER TOMÁS, M.D., DIMITRIO CARRIEDO, M.D., SANTIAGO MACÍAS, M.D., AND JESÚS BLANCO, M.D., FOR THE SPANISH LUNG FAILURE COLLABORATIVE GROUP*

VS et modes

Le Niveau d'Aide

- ***En fonction du VTE 6 à 8 ml.kg PBW***
- ***En fonction de l'effort: Pocc, P 0,1t, Pmo, EAdi Index Pdi, PTP oeso, Pmus***

Le bon niveau d'aide dans le sevrage ?

Determination of the optimal pressure support level evaluated by measuring transdiaphragmatic pressure.

T Kimura, J Takezawa, K Nishiwaki and Y Shimada

Chest 1991;100;112-117

Niveau d'aide ?

RESEARCH

Open Access

A novel non-invasive method to detect excessively high respiratory effort and dynamic transpulmonary driving pressure during mechanical ventilation

Michele Bertoni^{1,2}, Irene Telias^{3,4}, Martin Umer^{1,5}, Michael Long⁶, Lorenzo Del Sorbo^{3,2}, Eddy Fan^{1,2,7}, Christer Sinderby^{3,4}, Jennifer Beck^{3,4}, Ling Liu⁸, Haibo Qiu⁸, Jenna Wong², Arthur S. Slutsky^{3,4}, Niall D. Ferguson^{1,5,7,8,10}, Laurent J. Brochard^{3,4} and Ewan C. Goligher^{1,5,11,11*}

Computations

$$\text{Predicted } P_{mus} = -3/4 \times \Delta P_{occ}$$

$$\text{Predicted } \Delta P_L = (\text{Peak airway pressure} - \text{PEEP}) - 2/3 \times \Delta P_{occ}$$

Note: ΔP_{occ} is always ≤ 0 cm H₂O (the magnitude of decrease in airway pressure from inspiratory effort during the occlusion)

NARRATIVE REVIEW

Respiratory drive in the acute respiratory distress syndrome: pathophysiology, monitoring, and therapeutic interventions

Elena Spinelli¹, Tommaso Mauri^{1,2*}, Jeremy R. Beitler³, Antonio Pesenti^{1,2} and Daniel Brodie³

Driving Pressure Is Associated with Outcome during Assisted Ventilation in Acute Respiratory Distress Syndrome

Giacomo Bellani, M.D., Ph.D., Alice Grass, M.D.,
 Simone Gosio, M.D., Stefania Gatti, M.D.,
 Brian P. Kavanagh, M.B., Antonio Pesenti, M.D.,
 Giuseppe Foti, M.D.

ANESTHESIOLOGY 2019; 131:594–604

Pression Motrice en VSAI

LA SYNCHRONIE PATIENT VENTILATEUR

La synchronie

Impact of Expiratory Trigger Setting on Delayed Cycling and Inspiratory Muscle Workload

Didier Tassaux, Marc Gannier, Anne Battisti, and Philippe Joliet

Am J Respir Crit Care Med Vol 172. pp 1283–1289, 2005

Intensive Care Med
DOI 10.1007/s00134-005-0000-0

ORIGINAL

Udo Pfeiffer
Laurence Vignaux
Eric Baki
Jean Resche
Thierry Siffert
Noris-Emmanouil Lianos
Philippe Joliet
Didier Tassaux

Neurally adjusted ventilatory assist improves patient-ventilator interaction

Zoom choisi 00:00:10 (HH:MM:SS)

Détails des Zones d'intérêts

Nombre et Description

Durée de l'enregistrement 00:09:36:76
Fréquence : 25 Hz (HH:MM:SS:000)

VSAI: AI 0 ET PEP 0

Valve du respirateur ouverte, pas d'asservissement spirométrie et EAdi

Cyclage expiratoire

L'aide inspiratoire crée de l'asynchronie

Problématiques des asynchronies?

Intensive Care Med (2006) 32:1515–1522
DOI 10.1107/00134-006-0301-8

ORIGINAL

Arnaud W. Thille
Pablo Rodriguez
Belen Cabello
François Lellouche
Laurent Brochard

Patient-ventilator asynchrony during assisted mechanical ventilation

AI/VS PEP

Admettre patient

Nébuliseur

Etat

14/10

Pcrête (cmH ₂ O)	70
17	
PEP (cmH ₂ O)	4
F resp. (resp/min)	40
18	5
O ₂ (%)	40
Ti/Ttot	0.27
VMe (l/min)	20.0
C 9.6	5.0
Vc insp. (ml)	451
Vc exp. (ml)	453
Edi max (µV)	3.0
Edi min (µV)	0.1

Autres réglages

Conc. d'O₂
40
 21 % 100 0

PEP
7
 cmH₂O 50

Arrêt cycle
60
 % 70

Niv. AI sur PEP
12
 cmH₂O 120

Autres valeurs

Impact of Expiratory Trigger Setting on Delayed Cycling and Inspiratory Muscle Workload

Didier Tassaux, Marc Gannier, Anne Battisti, and Philippe Joliet

Am J Respir Crit Care Med Vol 172. pp 1283–1289, 2005

% débit de pointe ET 10

ET 25

ET 50

ET 70

AI/VS PEP

Admettre patient

Nébuliseur

Etat

14/10

Pcrête (cmH ₂ O)	19	70
PEP (cmH ₂ O)	6	
F resp. (resp/min)	21	5
O ₂ (%)	40	
Ti/Ttot	0.26	
VMe (l/min)	9.2	20.0
Vc insp. (ml)	445	5.0
Vc exp. (ml)	507	
Edi max (μV)	4.3	
Edi min (μV)	0.2	

Autres réglages

Conc. d'O₂ **40** %

PEP **7** cmH₂O

Arrêt cycle **60** %

Niv. AI sur PEP **12** cmH₂O

Autres valeurs

Arnaud W. Thille
Belen Cabello
Fabrice Galia
Aissam Lyazidi
Laurent Brochard

Reduction of patient-ventilator asynchrony by reducing tidal volume during pressure-support ventilation

Systeme Smartcare

- Systeme Néoganesch: règles, scénarios pour être dans une zone de confort: VTE FR E_TCO₂ évaluation 2-3 min

The knowledge base - 112 rules

SOME GENERAL RULES USED IN THE KNOWLEDGE-BASED SYSTEM

1. Never reduce the level of pressure support at night, except in case of excessive alveolar ventilation.
2. Try to keep the patient within a zone of "comfort," with a respiratory rate between 12 and 28 breaths per minute, a tidal volume above a minimum threshold, and a P_{ETCO2} below a maximum threshold.
3. Do not let the tidal volume fall below 300 or 250 ml in patients whose body weight is above or below 55 kg, respectively.
4. Keep the respiratory rate between 12 and 28 breaths/min so that the patient is comfortable. In some patients, the upper limit can be moved up to 32 breaths/min.
5. Do not let the end-tidal CO₂ exceed 60 mm Hg in COPD patients and 50 mm Hg in patients with other disorders.
6. Decrease the level of pressure support by 2 cm H₂O when the patient has a stable ventilation within the comfort zone during at least 30 min with a level of pressure support less than 20 cm H₂O.
7. Decrease the level of pressure support by 4 cm H₂O when the patient has a stable ventilation within the comfort zone during at least 60 min with a level of pressure support of 20 cm H₂O or more.
8. Initiate the preweaning observation period when the level of pressure support is at the minimal value (5 or 9 cm H₂O in tracheostomized or intubated patients, respectively).
9. Consider that the patient is ready to be weaned after 1 or 2 h of stable ventilation at the minimal level of pressure support (1 h in those patients with a level of pressure support of 15 cm H₂O or less after 1 h of observation, 2 h in those with an initial level of pressure support greater than 15 cm H₂O).
10. Adapt the level of pressure support to the physiologic needs of the patient and evaluate every 2 min.
11. Consider that a patient requiring a PEEP level above 5 cm H₂O is not ready to be weaned.
12. The maximal level of pressure support is 40 cm H₂O.
13. In case of severe hypoventilation, switch to assist-control ventilation with preset parameters.

A Multicenter Randomized Trial of Computer-driven Protocolized Weaning from Mechanical Ventilation

François Lellouche, Jordi Mancebo, Philippe Jolliet, Jean Roeseler, Frédérique Schortgen, Michel Dojat, Belen Cabello, Lila Bouadma, Pablo Rodriguez, Salvatore Maggiore, Marc Reynaert, Stefan Mersmann, and Laurent Brochard

Am J Respir Crit Care Med Vol 174. pp 894–900, 2006

Louise Rose
Jeffrey J. Presneill
Linda Johnston
John F. Cade

A randomised, controlled trial of conventional versus automated weaning from mechanical ventilation using SmartCare™/PS

SEPARATION DU VENTILATEUR

- ***INDEX PREDICTIFS***
- ***TEST***

The New England Journal of Medicine

©Copyright, 1991, by the Massachusetts Medical Society

Volume 324

MAY 23, 1991

Number 21

A PROSPECTIVE STUDY OF INDEXES PREDICTING THE OUTCOME OF TRIALS OF WEANING FROM MECHANICAL VENTILATION

KARL L. YANG, M.D., AND MARTIN J. TOBIN, M.D.

Premier Index FR/VT

Table 3. Effect of Duration of Mechanical Ventilation (<8 Days or ≥8 Days) on the Accuracy of Indexes in Predicting Weaning Outcome.*

INDEX	SENSITIVITY		SPECIFICITY		POSITIVE PREDICTIVE VALUE		NEGATIVE PREDICTIVE VALUE	
	<8 DAYS	≥8 DAYS	<8 DAYS	≥8 DAYS	<8 DAYS	≥8 DAYS	<8 DAYS	≥8 DAYS
Minute ventilation	0.79	0.75	0.75	0.08	0.65	0.35	0.40	0.33
Respiratory frequency	0.89	1.00	0.31	0.42	0.69	0.53	0.63	1.00
Tidal volume	1.00	0.88	0.50	0.58	0.78	0.58	1.00	0.88
Tidal volume/patient's weight	0.96	0.88	0.38	0.42	0.73	0.50	0.86	0.83
Maximal inspiratory pressure	1.00	1.00	0.00	0.25	0.64	0.47	1.00	1.00
Dynamic compliance	0.75	0.63	0.69	0.25	0.81	0.36	0.61	0.50
Static compliance	0.82	0.50	0.56	0.08	0.77	0.27	0.64	0.20
PaO ₂ /PAO ₂ ratio	0.79	0.88	0.38	0.17	0.69	0.41	0.50	0.67
Frequency/tidal volume ratio	1.00	0.88	0.63	0.67	0.82	0.64	1.00	0.89
CROP index	0.82	0.75	0.56	0.58	0.77	0.55	0.64	0.78

Identifying patients capable of breathing spontaneously and duration of mechanical ventilation

- **Intervention Group**
 - 1) **A daily screening of respiratory function (by the respiratory therapists of the unit)**
 - $\text{PaO}_2/\text{FIO}_2 > 200$
 - $\text{PEEP} < 5 \text{ cm H}_2\text{O}$
 - Adequate cough
 - $f/\text{VT} < 105 \text{ c/min}$
 - No vasopressor agents or sedatives
 - 2) **A 2-hour trial of spontaneous breathing**
 - 3) **Notification of the physician of the successful results**

Patients Receiving Mechanical Ventilation (%)

FEUX VERT

Neuro-ventilatory efficiency during weaning from mechanical ventilation using neurally adjusted ventilatory assist

H. Rozé^{1,2,3*}, B. Repousseau¹, V. Perrier¹, A. Germain¹, R. Séramondi¹, A. Dewitte^{1,3}, C. Fleureau¹ and A. Ouattara^{1,2}

Augmentation de l'EAdi pour un même niveau de pression entre le début et la fin du sevrage

TEST VSAI 7/0

Effect of Pressure Support vs T-Piece Ventilation Strategies During Spontaneous Breathing Trials on Successful Extubation Among Patients Receiving Mechanical Ventilation: A Randomized Clinical Trial

Carles Subirà, MD; Gonzalo Hernández, MD, PhD; Antònia Vázquez, MD, PhD; Raquel Rodríguez-García, MD; Alejandro González-Castro, MD; Carolina García, MD; Olga Rubio, MD, PhD; Lara Ventura, MD; Alexandra López, MD; María-Carmen de la Torre, MD, PhD; Eleni Keough, MD; Vanesa Arauz, MD; Carola Heemova, MD; Carmen Sánchez, MD; Ana Torán, MD; Iva Tejera, MD, PhD; César Lahera, MD; Sara Cabañes, MD; Victoria Lacueva, MD; María del Mar Fernández, MD, PhD; Anna Arnau, MSc, PhD; Rafael Fernández, RMD, PhD

USAI 7/0

Tube en T

Physiologic Effects of Reconnection to the Ventilator for 1 Hour Following a Successful Spontaneous Breathing Trial

Rémi Courmy, PhD; Alice Lejars, MD; Maeva Rodriguez, MD; Jean-Pierre Frat, PhD; Christophe Rault, PhD; François Arbidé, MD; Sylvain Le Pape, PhD; and Arnaud W. Thille, PhD

Noninvasive Positive-Pressure Ventilation for Postextubation Respiratory Distress

A Randomized Controlled Trial

Sean P. Keenan, MD, FRCPC, MSc

Caroline Powers, RRT

David C. McCormack, MD, FRCPC

Gary Block, MD, FRCPC

Outcomes	NPPV (n = 39)	Standard Therapy (n = 42)	P Value
Reintubation, No. (%)	28 (72)	29 (69)	.79
Pneumonia, No. (%)	16 (41)	17 (40)	.61
Duration of ventilation†			
Mean (SD)	8.4 (7.4)	17.5 (28.0)	.11
Median (range)	6.7 (0.5-28.6)	8.9 (2.0-146.7)	.12
ICU length of stay			
Mean (SD)	15.1 (10.9)	19.4 (25.0)	.32
Median (range)	11.9 (3.6-41.7)	10.8 (2.3-152.7)	.72
Hospital length of stay			
Mean (SD)	32.2 (25.4)	29.8 (28.4)	.69
Median (range)	19 (6-111)	22 (4-162)	.51
ICU survival, No. (%)	33 (85)	32 (76)	.34
Hospital survival, No. (%)	27 (69)	29 (69)	.99

ORIGINAL ARTICLE

Noninvasive Positive-Pressure Ventilation for Respiratory Failure after Extubation

Andrés Esteban, M.D., Ph.D., Fernando Frutos-Vivar, M.D.,
Niall D. Ferguson, M.D., Yaseen Arabi, M.D.,
Carlos Apezteguía, M.D., Marco González, M.D., Scott K. Epstein, M.D.,
Nicholas S. Hill, M.D., Stefano Nava, M.D., Marco-Antonio Soares, M.D.,
Gabriel D'Empaire, M.D., Inmaculada Alía, M.D., and Antonio Anzueto, M.D.

CONCLUSIONS

Noninvasive positive-pressure ventilation does not prevent the need for reintubation or reduce mortality in unselected patients who have respiratory failure after extubation.

Effect of Postextubation High-Flow Nasal Oxygen With Noninvasive Ventilation vs High-Flow Nasal Oxygen Alone on Reintubation Among Patients at High Risk of Extubation Failure A Randomized Clinical Trial

Sélection de patients à risque d'échec

Arnaud W. Thille, MD, PhD; Grégoire Muller, MD; Arnaud Gacouin, MD; Rémi Coudroy, MD; Maxens Decavèle, MD; Romain Schmeville, MD, PhD; François Bekmeier, MD; Christophe Girault, MD; Laurence Dangre, MD; Alexandre Lantier, MD, PhD; Sébastien Colasse, MD; Anahita Bouwé, MD; Emmanuel Vuze, MD; Anthony Le Meur, MD; Jean-Jamien Ricard, MD, PhD; Keyvan Razavi, MD; Guillaume Harbarth, MD; Christine Lebert, MD; Stephan Ehmann, MD, PhD; Caroline Sabatier, MD; Jeremy Bourenne, MD; Gaël Pradel, MD; Pierre Bailly, MD; Nicolas Terzi, MD, PhD; Jean Dellamonica, MD, PhD; Guillaume Lacave, MD; Pierre-Éric Danin, MD; Hodanou Nanadoumgar, MD; Aude Gibelin, MD; Lassane Zanre, MD; Nicolas Deye, MD, PhD; Alexandre Demoule, MD, PhD; Ariel Mazarin, MD; Mai Anh Ngy, MD; René Robert, MD, PhD; Stéphanie Bagot, PharmD, PhD; Jean-Pierre Frat, MD; for the HIGH-WEAN Study Group and the REVA Research Network

No. at risk									
High-flow nasal oxygen									
Alone	302	276	265	253	248	246	244	243	
With noninvasive ventilation	339	321	314	308	305	294	292	291	

Effect of Protocolized Weaning With Early Extubation to Noninvasive Ventilation vs Invasive Weaning on Time to Liberation From Mechanical Ventilation Among Patients With Respiratory Failure The Breathe Randomized Clinical Trial

Pas de précipitation

Gavin D. Perkins, MD; Dipesh Mistry, PhD; Simon Gates, PhD; Fang Gao, MD; Catherine Snelson, MD; Nicholas Hart, PhD; Luigi Camporota, PhD; James Varley, MB; Coralia Carle, MB; Flankumaran Paramasham, MB; Rowley Hoddrell; Daniel T. McAuley, MD; Timothy S. Walsh, MD; Bronagh Blackwood, PhD; Louise Rose, PhD; Sarah F. Lamb, DPhil; Stavros Petrou, PhD; Duncan Young, DM; Ranjit Lall, PhD; for the Breathe Collaborators

Conclusion

- Protocole de sevrage
- Screening quotidien des patients
- Adapter le niveau d'aide
- Améliorer l'interaction patient ventilateur
 - Synchronisation
 - Proportionnalité
 - Nouvel outil de monitoring au lit du patient: l'EAdi