



Trachéotomie en Réanimation



DU Kiné
Mars 2026



Pr Hadrien Rozé
Réanimation Polyvalente
Centre Hospitalier Côte Basque

www.slm-va.com

JAMA Otolaryngology-Head & Neck Surgery | Original Investigation

Timing, Complications, and Safety of Tracheotomy in Critically Ill Patients With COVID-19

Francisco Xavier Avilés-Jurado, MD, PhD; Daniel Prieto-Alhambra, MD, PhD; Neely González-Sánchez, MD; José de Ossa, MD; Claudio Arancibia, MD; María Jesús Rojas-Lechuga, MD; Laura Ruiz-Sevilla, MD; Joan Remacha, MD; Irene Sánchez, MD; Eduardo Lehrer-Coriat, MD; Mauricio López-Chacón, MD; Cristóbal Langdon, MD; Josep Maria Gallemany, MD, PhD; Francisco Larrosa, MD, PhD; Isam Alkoidi, MD, PhD; Manuel Bernal-Sprekelsen, MD, PhD; Pedro Castro, MD, PhD; Isabel Vilaseca, MD, PhD


JAMA Otolaryngol Head Neck Surg. 2021;147(10):41-48.

Characteristic	Total (N=50)	Early tracheotomy (n=20)	Late tracheotomy (n=30)
Postoperative complications			
Minor diffuse bleeding	6 (12)	5 (15)	1 (3)
Major bleeding	0	0	0
Air leak	3 (6)	3 (15)	0
Cardiovascular	1 (2)	0	1 (3)
Tolerance to cuff deflation in survivors (n = 42)	42 (100)	42 (100)	42 (100)
Ability to phonate in survivors (n = 42)	42 (100)	42 (100)	42 (100)
Neurologic impairment			
No	41 (82)	28 (56)	13 (26)
Yes	9 (18)	2 (4)	7 (14)
Death before extubation	6 (12)	2 (4)	4 (8)
Intubation time before tracheotomy, d			
Median (IQR)	9 (4-25)	8 (2)	12.5 (2-25)
Mean (SD)	9.6 (9.4)	7.5 (2.9)	12.2 (9)
Total time receiving IMV, d			
Median (IQR)	19 (6)	17 (5)	20 (5)
Mean (SD)	17.9 (4.5)	16.6 (4.5)	20.5 (3.4)
Time from tracheotomy to IMV withdrawal, d			
Median (IQR)	8 (6)	9 (6)	8 (6)
Mean (SD)	8.5 (4.5)	9.2 (5.1)	7.2 (2.8)
Decannulation rate in survivors	42 (100)	42 (100)	42 (100)
Time to decannulation, d			
Median (IQR)	22 (15)	20 (15)	25 (14)
Mean (SD)	23.7 (11)	22.2 (13)	26.4 (9)

• Equation du mouvement du poumon :

$$P_{alv} = PEP + \frac{V}{Ctp} + (Débit \times Res)$$

RS vs VC
VNI vs IOT
VAS nez vs bouche
Trachéo vs nez




Labels: Cage thoracique, Cotes, Action de la cage thoracique, Trachée, Plèvre viscérale, Poumon, Plèvre pariétal, Pressure, PIP, Pplat, PEP, time.

Propriétés résistives

- Ventilation = condition dynamique
- L'activité des muscles respiratoires doit
 - vaincre l'élasticité pulmonaire (2/3 au repos)
 - vaincre la résistance du système respiratoire au
- passage de l'air (1/3 au repos)
 - résistances tissulaires (≈ 20%): frottements du tissu pulmonaire
- résistances des voies aériennes (≈ 80%): résistance à l'écoulement des molécules

• Fluide

- viscosité η
- densité ρ
- conduit de longueur l
- rayon r




• En écoulement laminaire

$$\dot{V} = \frac{\Delta P^*}{R} \quad R = \frac{8\eta l}{\pi r^4}$$

Résistances pulmonaires

En respiration nasale



- VA extra-thoraciques (nez +++)
50%
- Trachée et grosses bronches
40%
- G_7 à G_{23}
10%

Resistances Trachéo = VS par le nez

- R trac = R VS par le nez
- Si IOT la longueur modifie les résistances inspi et expiratoires et R Trac < R IOT à même diamètre (7 ou 8)



Espace mort



- L'augmentation de l'effet espace mort (de l'ordre de 25 % chez le sujet normal et pouvant atteindre 60 à 80 % chez le patient BPCO en poussée) est responsable de l'hypercapnie.
- La PaCO₂ est proportionnelle à la production de CO₂ et inversement proportionnelle à la ventilation alvéolaire (VA) :
- PaCO₂ = K x VCO₂/VA
- La ventilation alvéolaire est la différence entre la ventilation totale et la ventilation de l'espace mort :
- VA = Ve x VD/VT = VE x (1-VD)/VT
- VD est l'espace mort physiologique, c'est-à-dire la fraction du volume courant qui ne participe pas aux échanges gazeux. VT est le volume courant. Ainsi, en remplaçant VA dans la première équation, on obtient :
- PaCO₂ = K x VCO₂ / (VE x (1-VD)/VT)

Espace mort

- La trachéo diminue l'espace mort de 156 ml à 230 ml.
- Le VT est poché de 6 ml.kg afin de limiter les asynchronies et l'auto PEP chez les BPCO



Intensive Care Med (2002) 28:1761-1767


Intensive Care Med (2008) 34:1477-1486



$$P_{tot(t)} = R \cdot \text{Débit}(t) + 1/C$$

$$P_{tot} = P_{mus} + P_{aw}$$





- Assistance en fonction du travail respiratoire

Intensive Care Med (2006) 32:1311-1314
DOI 10.1007/s00134-006-0278-3 PHYSIOLOGICAL NOTE

1 joule = mobilisation de 1l de gaz avec un ΔP de 10 cmH₂O
En (joule) $\neq P$

Belen Cabello
Jordi Mancho


Work of breathing

Review First published April 30, 2006, doi:10.1155/2006/40003000

HIGHLIGHTED TOPIC | The Respiratory Muscles in Chronic Obstructive Pulmonary Disease

Role of the respiratory muscles in acute respiratory failure of COPD: lessons from weaning failure

Martin J. Tobin, Franca Leblak and Laurent Brochard




WOB: Travail respiratoire

- 31 BPCO échec de sevrage
- Début SBT: PTP échec 255 (59) cmH₂O.s/min succès 158 (23)
- Fin SBT : PTP échec 388 (68)* succès 205 (25)

>> Echec si trop de travail

Jubran, Tobin. Am J Respir Crit Care Med 1997;155: 906-915



AVANTAGES ATTENDUS

REDUCTION DU TRAVAIL RESPIRATOIRE




DIMINUTION DES PNEUMOPATHIES

SEVRAGE VENTILATOIRE PROGRESSIF

CONFORT

SECURITE

© IST medical





Impact of tracheotomy on sedative administration, sedation level, and comfort of mechanically ventilated intensive care unit patients*


Ania Nieszkowska, MD; Alain Combes, MD, PhD; Charles-Edouard Luyt, MD, PhD; Hichem Ksibi, MD; Jean-Louis Trouillet, MD; Claude Gibert, MD; Jean Chastre, MD

Characteristic	72 patients avant vs après trachéo	Value
At ICU admission		
Age		58 (52,72)
Sex, male, n (%)		43 (59.7)
Type of admission, n (%)		
Surgery		40 (55.6)
Cardiac		38 (52.7)
Other		2 (2.8)
Medical		32 (44.4)
SAPS II		46 (36,66)
McCabe		2 (1,2)
Reason for MV, n (%)		
Acute exacerbation of COPD		7 (9,7)
Acute respiratory failure		29 (40,3)
Postoperative respiratory failure		27 (37,5)
Neurologic		9 (12,5)
On the day of tracheotomy		
SOPA score		7 (5,8,10,0)
Radiologic score		5 (3,7)
Temperature, °C, mean ± SD		37,8 ± 1,0
PaO ₂ /P _{iO₂} , mm Hg		197 (150,280)
Shock, n (%)		29 (40,3)

Nieszkowska Crit Care Med 2005



Nutrition voie orale




Nieszkowska Crit Care Med 2005

Outcome of mechanically ventilated patients who require a tracheostomy*


Fernando Frutos-Vivar, MD; Andrés Esteban, MD, PhD; Carlos Apezteguía, MD; Antonio Anzueto, MD; Peter Nightingale, MD; Marco González, MD; Luis Soto, MD; Carlos Rodrigo, MD; Jean Raad, MD; Cide M. David, MD; Dimitros Matamis, MD; Gabriel D'Empaire, MD; for the International Mechanical Ventilation Study Group

- Mars 1998, 361 services, 20 pays
- 5 183 patients ventilés > 12 h (- 102 déjà trachéotomisés)
- 546 (10,7 %) trachéotomisés après 12 [7-17] j de VM

CCM 2005;33:290-298



CCM 2005;33:290-298



CCM 2005;33:290-298




Figure 3. Tree-building of risk factors for tracheostomy obtained by the recursive partitioning method.

CCM 2005;33:290-298

Trachéotomisés : Quel devenir ?

	Mortalité en réa	Mortalité hospitalière
Trachéotomie +	20 %	39 %
Trachéotomie -	32 %	40 %

CCM 2005;33:290-298

Tracheostomy does not improve the outcome of patients requiring prolonged mechanical ventilation: A propensity analysis*

Christophe Clec'h, MD; Corinne Alberti, MD, PhD; François Vincent, MD; Maïté Garrouste-Orgeas, MD; Arnaud de Lassence, MD; Dany Toledano, MD; Elie Azoulay, MD, PhD; Christophe Adrie, MD; Samir Jamali, MD; Isabelle Zaccaria; Yves Cohen, MD; Jean-François Timsit, MD, PhD; on behalf of the OUTCOMEREA study group

Critical Care Medicine

Patients comparables

Variable	Tracheostomy (n = 160)	No Tracheostomy (n = 422)	p Value
Duration of MV ≥15 days, no. (%)	95 (59.4)	229 (54.3)	.77
Need for reintubation, no. (%)	4 (2.5)	6 (1.3)	.25
MV for neurologic disease, no. (%)	46 (28.8)	126 (27.8)	.67
Chronic respiratory disease, no. (%)	46 (28.8)	101 (22.3)	.11
Do-not-resuscitate order, no. (%)	18 (11.3)	61 (13.4)	.48
Cause of acute respiratory failure, no. (%)			
Postoperative	24 (15)	82 (18.1)	.16
Congestive heart failure	10 (6.3)	32 (7.1)	.99
Cardiac arrest	4 (2.5)	8 (1.8)	.46

CCM 2007;35:132-138

Mortalité en réanimation

	OR	95% CI	p Value
Model 1			
All patients	0.94	0.63-1.39	.74
Patients with early tracheostomy ^a	0.41	0.10-1.80	.24
Patients with late tracheostomy ^b	0.97	0.65-1.50	.90
Model 2			
All patients	1.12	0.75-1.67	.59
Patients with early tracheostomy ^a	0.78	0.21-2.91	.71
Patients with late tracheostomy ^b	1.16	0.77-1.75	.49

CCM 2007;35:132-138

Effect of Early vs Late Tracheostomy Placement on Survival in Patients Receiving Mechanical Ventilation

The TracMan Randomized Trial

JAMA. 2013;309(20):2121-2129

Early, days 1-4

Late, day 10 or later if clinically indicated

No. of patients

Days From ICU Admission to Tracheostomy

Effect of Early vs Late Tracheostomy Placement on Survival in Patients Receiving Mechanical Ventilation

The TracMan Randomized Trial

JAMA. 2013;309(20):2121-2129

Cumulative Survival

Years After Randomization

No. at risk	0	0.5	1.0	1.5	2.0
Early	451	261	244	230	221
Late	448	242	226	217	205

TRACHÉOTOMIE PRÉCOCE VERSUS TRACHÉOTOMIE TARDIVE / INTUBATION PROLONGÉE

Trachéotomie précoce: < 8 jours d' intubation
Trachéotomie tardive: > 15 jours d' intubation

Méta-analyse de Griffiths BMJ 2005
Méta-analyse de Wang Chest 2010

Systematic review and meta-analysis of studies of the timing of tracheostomy in adult patients undergoing artificial ventilation

John Griffiths, Vicki S Barber, Lesley Morgan, J Duncan Young

Study	No of patients (n/N)	Timing of tracheostomy		Intensive care setting	Randomisation	Mortality expressed on intention to treat basis	Duration of ventilation and critical care site expressed on intention to treat basis
		Early	Late				
Boudier et al 2004 ²⁷	62	5-6 days after admission	Prolonged endotracheal intubation	Unit for patients with head injuries	Randomised; method not stated	Implied	Implied both
Dunham et al 1984 ²⁸	74	3-4 days after initiation of transaryngeal intubation	14 days after initiation of transaryngeal intubation	Trauma unit	Quasi-randomised	Mortality not recorded	Yes
Rodriguez et al 1990 ²⁹	106	1-7 days after admission to intensive care unit	8 or more days after admission to intensive care unit	Surgical unit	Quasi-randomised	Implied	Implied both
Rumbak et al 2004 ³⁰	120	0-7 days after initiation of mechanical ventilation	14-16 days after initiation of mechanical ventilation	Three medical units	True randomisation	Implied	Yes
Saffie et al 2002 ³¹	44	Next available operative day	14 days after burn injury	Burns unit	True randomisation	Implied	Yes


2005 BMJ Griffiths

Systematic review and meta-analysis of studies of the timing of tracheostomy in adult patients undergoing artificial ventilation

John Griffiths, Vicki S Barber, Lesley Morgan, J Duncan Young



Mortalité en réanimation




2005 BMJ Griffiths

Systematic review and meta-analysis of studies of the timing of tracheostomy in adult patients undergoing artificial ventilation

John Griffiths, Vicki S Barber, Lesley Morgan, J Duncan Young




Pneumonies infectieuses



2005 BMJ Griffiths

Durées de ventilation



2005 BMJ Griffiths

CHEST Original Research

The Timing of Tracheostomy in Critically Ill Patients Undergoing Mechanical Ventilation

A Systematic Review and Meta-analysis of Randomized Controlled Trials

Fei Wang, MD, PhD; Yongping Wu, MD, PhD; Luoguo Bo, MD, PhD; Junghong Lee, MD, PhD; Jiahui Zhu, MD; Feng Chen, MD, PhD; Jiahao Li, MD, PhD; and Xiaoming Deng, MD, PhD


Table 1. Characteristics of Included Trials

Study/Year	Randomized Controlled Trial	Timing of Tracheostomy	Mean Age (y)	Duration of Mechanical Ventilation (days)	Exclusion Criteria	Study Quality	Study Location
Saffie et al 2002 ¹⁶	Yes	Early vs Late	44.5 (19.7)	52.1 (19.2)	Not reported	Yes	China
Bouderka et al 2004 ²⁴	Yes	Early vs Late	41.1 (17.4)	40 (19)	Not reported	Yes	China
Rodriguez et al 1990 ²⁶	Yes	Early vs Late	40 (15.3)	40 (15.3)	Not reported	Yes	Spain
Rumbak et al 2004 ¹⁷	Yes	Early vs Late	57.7 (13.3)	40 (15.3)	Not reported	Yes	China
Terragni et al 2010	Yes	Early vs Late	58 (10.8)	58 (10.8)	Not reported	Yes	Italy
Trouillet et al 2011	Yes	Early vs Late	61.3 (16.4)	61.3 (16.4)	Not reported	Yes	France

CHEST 2011; 140(6):1456-1463


DURÉE DE VENTILATION

(trachéotomie précoce versus tardive)




Wang Chest 2011



influence sur la mortalité précoce



influence sur la mortalité tardive



Wang Chest 2011



Annals of Internal Medicine | ORIGINAL RESEARCH

Early Percutaneous Tracheotomy Versus Prolonged Intubation of Mechanically Ventilated Patients After Cardiac Surgery

A Randomized Trial*

Jean-Louis Trouillet, MD; Charles-Edouard Loyt, MD, PhD; Marguerite Guiguet, PhD; Alexandre Ouattara, MD, PhD; Elisabeth Vaissier, MD; Ralouka Makri, MD; Ania Nieszewska, MD; Pascal Leprince, MD, PhD; Alain Pavie, MD; Jean Chastre, MD; and Alain Combes, MD, PhD

Sedation	ET (n=177)	LT (n=206)	Mean Difference (95% CI)	P Value
Mean duration of intravenous sedation (SD), d†	6.4 (5.9)	9.6 (7.3)	-3.2 (-5.0 to -1.3)	0.007
Mean sedation-free days during 1-28 d (SD)	19.0 (9.1)	15.5 (9.3)	4.5 (1.2 to 6.9)	0.005
Mean cumulative sufentanil dose during 1-15 d (SD), µg/kg	4.0 (6.5)	10.2 (18.2)	-6.2 (-9.9 to -2.5)	0.001
Mean cumulative propofol dose during 1-15 d (SD), mg/kg	32.9 (60.2)	67.8 (116.7)	-34.9 (-60.1 to -9.8)	0.004
Mean cumulative midazolam dose during 1-15 d (SD), mg/kg	2.7 (4.7)	6.4 (14.3)	-3.7 (-6.6 to -0.8)	0.01
Mean days (during 1-15 d) of haloperidol therapy (SD)	1.9 (3.0)	3.2 (4.2)	-1.3 (-2.3 to -0.3)	0.01
Mean cumulative haloperidol dose during 1-15 d (SD), mg/kg	0.26 (0.51)	0.57 (0.92)	-0.3 (-0.5 to -0.1)	0.002
VAP after randomization, n (%)	50 (46)	47 (44)	2.0 (-1.3 to 5.2)	0.77
Sternal wound infection, n (%)	14 (13)	14 (13)	-0.2 (-3.2 to 2.8)	0.96
Bloodstream infection, n (%)	18 (17)	16 (15)	1.5 (-8.1 to 11.3)	0.85
Mean days (during 1-15 d) nurse-assessed as comfortable (SD)	11.8 (3.8)	10.4 (4.4)	1.4 (0.3 to 2.5)	0.01
Mean days (during 1-15 d) with nurse-assessed easy management (SD)	12.0 (3.8)	10.8 (4.4)	1.2 (0.09 to 2.3)	0.04
Received oral nutrition at 15 d, n (%)	91 (83)	57 (53)	30.2 (18.5 to 42.2)	<0.001
Bed-to-chair transfer at 15 d, n (%)	72 (66)	47 (44)	22.1 (9.2 to 35.1)	0.002
Muscle strength assessment (SD)				
14 d (n = 76, 68)	156.9 (87.0)	134.9 (92.8)	22.0 (-7.7 to 51.6)	0.15
28 d (n = 36, 30)	164.1 (86.1)	176.9 (85.6)	-12.9 (-53.3 to 27.5)	0.52
42 d (n = 21, 21)	170.1 (86.4)	159.4 (67.9)	10.7 (-13.6 to 25.1)	0.30
56 d (n = 8, 11)	149.7 (70.4)	185.4 (76.0)	-35.7 (-108.0 to 36.6)	0.31

Ann Intern Med. 2011;154:373-383.

Is the Duration of Mechanical Ventilation Predictable?*

Gilles Troché, MD; and Pierre Moine, MD

CHEST 1997; 112:745-51

- Réanimation chirurgicale
- 195 patients + cohorte de validation (n = 128)
- Prédiction d'une VM ≥ 15 j
- Multiples variables analysées : Age, BMI, type chirurgie, IGS, LIS, OSF, ...
- Régression logistique, ROC curve

Difficile à prévoir !

Parameters	Group 1 (MV≤14 d) (n=177)	Group 2 (MV≥15 d) (n=206)	P Value
Age, yr	59.7 ± 19.6	57.0 ± 20.6	NS (0.52)
BMI	24.5 ± 5.0	24.6 ± 4.2	NS (0.92)
Emergent admission	127 (71.8%)	23 (88.5%)	NS (0.07)
Elective admission	50 (28.2%)	3 (11.5%)	
Altecsier classification			NS (0.30)
0 (medical patients)	38 (21.5%)	7 (36.9%)	
I and II	83 (30.0%)	4 (15.4%)	
III and IV	86 (48.6%)	15 (57.7%)	
Pathology/admission			NS (0.31)
Abdominal surgery	106 (59.9%)	12 (46.2%)	
Orthopedic surgery	3 (1.7%)	6 (23.1%)	
Gynecologic or obstetric disease	6 (3.4%)	0	
Medical disease	42 (23.7%)	8 (30.8%)	
SAPS	15.1 ± 5.6	15.5 ± 4.2	NS (0.77)
CCS	13.0 ± 3.7	12.3 ± 3.6	NS (0.34)
APACHE II	20.7 ± 8.3	22.9 ± 7.7	NS (0.20)
SS	9.2 ± 5.7	12.5 ± 7.5	0.01
OSFI 0	95 (53.7%)	6 (23.1%)	
OSFI I	51 (28.8%)	14 (53.8%)	<0.05
OSFI II, III, or IV	31 (17.5%)	6 (23.1%)	
Albuminemia, g/L	38.1 ± 7.7	34.1 ± 7.5	<0.05

*NS = not significant.

Shock on Admission Day Is the Best Predictor of Prolonged Mechanical Ventilation in the ICU*

Elisa Estenssoro, MD; Francisco González, MD; Enrique Laffaire, MD; Héctor Canales, MD; Gabriela Sáenz, MD; Rosa Reina, MD; and Arnaldo Dubin, MD

- Réanimation polyvalente (Buenos Aires)
- 551 patients (mortalité 31 %), 348 ventilés
- Multiples variables recueillies à J 1
- Ventilation prolongée : > 21 j (n = 79, 23 % des ventilés)

CHEST 2005; 127:598-603

Shock on Admission Day Is the Best Predictor of Prolonged Mechanical Ventilation in the ICU*



Elisa Estenssoro, MD; Francisco González, MD; Enrique Laffaire, MD; Héctor Canales, MD; Gabriela Sáenz, MD; Rosa Reina, MD; and Arnaldo Dubin, MD

Variables	OR	95% CI	p Value
Shock day 1	2.17	1.25–3.77	0.006
SAPS II	1.04†	1.02–1.06	0.0001
PaO ₂ /F _i O ₂ day 1	0.99‡	0.99–1.00	0.125

CHEST 2005; 127:598–603

Quels patients trachéotomiser ?

Concept de « chronically ill patients »

- Survivants de situations dramatiques
- Dépendants Ventilation Mécanique
- Dépendants richesse environnement humain et technique
- Sans autre défaillance vitale
- Nombreuses autres affections : séquelle(s) des défaillances viscérales, motrice, déficit nutritionnel, germes nosocomiaux, processus invasifs, ...

⇒ Typologie particulière



Mobilizing Patients in the Intensive Care Unit Improving Neuromuscular Weakness and Physical Function



Dale M. Needham, MD, PhD



Sevrage et Réhabilitation : Décanulation

Intensive Care Med (2003) 29:845–848
DOI 10.1007/s00134-003-1689-z

BRIEF REPORT

Piero Ceriana
Annalisa Carfacci
Paolo Navalesi
Ciro Rampulla
Monica Delmastro
GianCarlo Piaggi
Elisa De Mattia
Stefano Nava

Weaning from tracheotomy in long-term mechanically ventilated patients: feasibility of a decisional flowchart and clinical outcome


« Pré requis »

- **Stabilité clinique**
- **PaCO₂ < 60 mmHg**
- **Absence de délire ou d'autres désordres psychiatriques**
- **Absence de sténose trachéale ou glottique**
- **Toux efficace**, pression expiratoire maximale (PEM) ≥ 40 cmH₂O
- **Déglutition correcte**
- **Patient consentant**

Respiratory Muscle Dysfunction Secondary to Chronic Tracheostomy Tube Placement*



Gerard Criner, M.D.,† Barry Make, M.D., F.C.C.P.; and Bartolome Celli, M.D., F.C.C.P.




	Tracheostomy†	Montgomery Button	Decannulation	Predicted
FVC (L)	1.17	0.88	1.02	2.94
FEV ₁ (L)	0.52	0.44	0.48	2.32
FEV ₁ /FVC	0.44	0.50	0.47	0.80
FRC (L)	4.50		3.99	2.62
Raw (cmH ₂ O/L/s)	7.55	4.28	2.94	2.29
Po ₂ (mm Hg)	80	80	N†	N†
Pco ₂ (mm Hg)	51	54	N‡	N‡

Intensive Care Med (2002) 28:1761–1767
DOI 10.1007/s00134-002-1545-6



ORIGINAL

Karim Chadda
Bruno Louis
Lamine Benabiss
Djillali Amrane
Philippe Gajdos
Jean Claude Raphael
Frédéric Lofaso

Physiological effects of decannulation in tracheostomized patients



$$V_D^{phys} + V_D^{circ} = V_T \times (P_{aCO_2} - P_{E_{CO_2}} / P_{aCO_2})$$



CONCLUSIONS

La trachéotomie n'est pas une technique de sevrage

La trachéotomie est bénéfique uniquement par rapport à la ventilation prolongée

La trachéotomie garde toute sa place pour les malades au sevrage très prolongé

Une stratégie visant à aller vers la décanulation est nécessaire

Des structures spécialisées peuvent être intéressantes